PUBLICATIONS OF THE UNIVERSITY OF MANCHESTER

MEDICAL SERIES
No. X.

Lectures on the Pathology of Cancer

SHERRATT & HUGHES
Publishers to the Victoria University of Manchester
Manchester: 34 Cross Street
London: 33 Soho Square, W.

Lectures on the Pathology of Cancer

BY

CHARLES POWELL WHITE, M.A., M.D., F.R.C.S.
Pilkington Cancer Research Fellow

MANCHESTER
AT THE UNIVERSITY PRESS
1908

2980

616·994
No 8

UNIVERSITY OF MANCHESTER PUBLICATIONS
No. XLII.

PREFACE.

HESE lectures were delivered in the University of anchester in June and July, 1908, in connection with e Pilkington Cancer Research Fund. They do not ofess to contain the whole pathology of Cancer. My ea has been rather to give a general outline, dealing ore particularly with certain points which are neglected, imperfectly dealt with, in most works on the subject. ence I have laid particular stress on the life history of mours, and on certain biological problems connected erewith.

I desire to take this opportunity of expressing my anks to Mrs. Pilkington, the founder of the Pilkington ancer Research Fund; and to Professor Lorrain Smith, whose department my work is carried on.

CHARLES POWELL WHITE.

ANCHESTER,
October, 1908.

CONTENTS.

PAGE

PART I. Tumours in General 3

Definition—Classification—Nomenclature—Relation of Tumour-growth to other Processes—Difference between Simple and Malignant Tumours—Rudiment of Origin—Mode of Growth.

PART II. Cancer 17

Cytomata—Cancer in Animals—Rudiment of Origin—Mode of Growth—Modes of Extension—Multiple Tumours—The Stroma of Carcinoma—Vascular Supply—Ultimate Destination —Degenerative Processes—Metabolism—Clinical Features.

PART III. Causation 33

Causal Factors—Extrinsic Causal Factors—Intrinsic Causal Factors—Cell Proliferation—Influences which limit Proliferation—Co-ordinating Mechanism of the Organism—The Relation of the Tumour to the Organism—The Nature of the Intercellular Influences—Physiological Equilibrium—The Influence of Inflammatory Conditions—The Influence of Heredity—The Influence of Civilisation—The Influence of Sex, Age, and Site.

PART IV. The Bearings of Pathology on Treatment ... 59

Recapitulation—Principles of Treatment—Conclusion.

APPENDIX. The Parasitic Theory 69

LIST OF ILLUSTRATIONS.

Fig. 1. Adenoma of the Liver.
Fig. 2. Carcinoma of the Liver.
Fig. 3. Chondroma.
Fig. 4. Chondrosarcoma.
Fig. 5. Adenoma of the Rectum.
Fig. 6. Carcinoma of the Rectum.
Fig. 7. Carcinoma of the Rectum.
Fig. 8. Carcinoma of the Rectum.
Fig. 9. Angeioma of the Liver.
Fig. 10. Blastocytoma of the Kidney.
Fig. 11. Blastocytoma of the Kidney.
Fig. 12. Blastocytoma of the Kidney.
Fig. 13. Blastocytoma of the Kidney.
Fig. 14. Blastocytoma of the Kidney.
Fig. 15. Blastocytoma of the Parotid.
Fig. 16. Primary Carcinoma of the Femur.
Fig. 17. Primary Carcinoma of the Pleura.
Fig. 18. Primary Carcinoma of a Rib.
Fig. 19. Sarcoma of the Neck from a Guinea Pig.
Fig. 20. Carcinoma of the Thyroid from a Mouse.
Fig. 21. Lipoma from a Parrot.
Fig. 22. Giant celled Sarcoma of the Femur from a Drake.
Fig. 23. Carcinoma of the Kidney from a Rabbit.
Fig. 24. Carcinoma of the Kidney from a Rabbit.
Fig. 25. Carcinoma of the Breast infiltrating muscle.
Fig. 26. Carcinoma of the Breast infiltrating fat.
Fig. 27. Carcinoma infiltrating a vein.
Fig. 28. Carcinoma of the Breast. Polarised light.
Fig. 29. Adrenal body. Polarised light.
Fig. 30. Adrenal body. Polarised light.
Fig. 31. Adrenal body. Polarised light.
Fig. 32. Bilharzial Carcinoma of the Bladder.
Fig. 33. Bilharzial Carcinoma of the Bladder.

PART I.

TUMOURS IN GENERAL

PART I. TUMOURS IN GENERAL.

DEFINITION.

To understand the pathology of Cancer it is necessary first to know what Cancer is, what its relationship is to other tumours, to other diseases, and to normal tissues.

By Cancer I understand any form of malignant disease. Some authors have attempted to limit the term to Carcinoma but there is no reason for this, and it is more satisfactory to use the compact term Cancer as a synonym for the more cumbrous term Malignant disease.

The cancers are members of the group of disorders known as Tumours which are characterised especially by their progressive and independent growth, and the pathology of Cancer cannot be considered apart from the pathology of other tumours.

The definition and classification of tumours which I adopt is one which I brought forward in the Erasmus Wilson lectures which I delivered before the Royal College of Surgeons in 1902.[10]

A tumour is defined as *a mass of cells, tissues, or organs, resembling those normally present in the body, but arranged atypically, which grows at the expense of the organism without, at the same time, subserving any useful function therein.*

There are two important points in this definition. First, the distinguishing character of a tumour is the fact that it is *atypical*, not in the structure, but in the arrangement of its component parts. This feature distinguishes a tumour from a condition such as hypertrophy. The second point is that a tumour grows at the expense of the organism, but does not subserve any useful function therein. This distinguishes a tumour from an inflammatory mass, which, however destructive it may appear, has a useful tendency and ceases to increase when the irritant which causes it is removed or neutralised.

The normal organism is composed of a number of *organs* of which the structure and arrangement are characteristic for the species but differ in different species.

An organ is composed of a combination of *tissues* of which the structure and arrangement are typical for the organ concerned but differ in different organs.

A tissue is composed of a number of *cells* with more or less intercellular substance, of which the structure and arrangement are typical for each tissue, but differ in different tissues.

If this typical arrangement fail at any point the result is a tumour of some kind.

Thus we may find *organs*, recognisable as such, arranged in an atypical manner as in a *dermoid cyst* in which we can recognise skin, teeth, glands, bones, etc., without any definite arrangement.

Also *fibrous tissue* with its blood-vessels, etc., is normally arranged to form a definite organ—a tendon or fascia. If, however, while retaining its typical characters as fibrous tissue, it is not arranged in a typical manner, but forms an irregular mass we have a tumour—a *fibroma*. Again *fibrous tissue cells* are normally arranged with a certain amount of intercellular substance to form a tissue—*fibrous tissue*. If, however, the cells multiply without forming the intercellular substance the result is a tumour—a *sarcoma*.

To take another example: *epithelium* is normally arranged in a definite manner together with connective-tissue, blood-vessels, etc., to form an organ, for example, a secreting gland. In an *adenoma,* however, the epithelium and connective-tissue, while themselves resembling normal tissues, are arranged irregularly and not so as to form a gland (Fig. 1). Again, normally *epithelial cells* are arranged in contact with one another forming a membrane which covers a surface or lines tubes and spaces. This membrane is an epithelium—a tissue. If, however, the epithelial cells are not arranged in the typical way but grow irregularly in the connective-tissue spaces not forming a definite membrane, we have a tumour—a *carcinoma* (Fig. 2).

CLASSIFICATION.

We have, then, the following classification of tumours.

In the first place we have tumours which are composed of more or less definite *organs* which are arranged atypically as in a dermoid. These are called ORGAN TUMOURS or ORGANOMATA.

In the next place we have tumours which are composed of *tissues* arranged atypically. These are called TISSUE TUMOURS or HISTIOMATA and include the non-malignant tumours.

Lastly we have tumours which are composed of *cells* arranged atypically. These are called CELL TUMOURS or CYTOMATA and include the cancers.

In this last group the characteristic feature is to be found in the growing portion of the tumour. Some carcinomata show well formed epithelium and closely resemble adenomata, and some sarcomata contain well formed fibrous tissue, cartilage, etc., but, if the growing part of the tumour be examined the distinction will be found to hold good, the definite tissues being of secondary formation whereas in the histiomata the definite tissues are present in the growing as in the older parts. I shall have more to say on this point when I consider the growth of tumours.

The classification is, then, as in the accompanying table.

CLASSIFICATION OF TUMOURS.

A. ORGANOMATA.	Teratoma. (Dermoid Cyst.)
B. HISTIOMATA.	
(*a*) Connective tissue tumours.	DESMOMATA.
Fibrous tissue,	Fibroma.
Fat,	Lipoma.
Mucous tissue,	Myxoma.
Notochordal tissue,	Chordoma.
Cartilage,	Chondroma.
Bone,	Osteoma.
Neuroglia,	Glioma.
(*b*) Lymphoid tissue tumours.	LYMPHOMATA.
Lymphadenoid tissue,	Lymphadenoma. (Lymphoma.)
Bone marrow,	Myeloma.
(*c*) Muscle tumours.	MYOMATA.
Smooth muscle,	Leiomyoma.
Striated muscle,	Rhabdomyoma.
(*d*) Nerve tissue tumours.	NEUROMATA.
Medullated Nerve tissue,	Myelinic Neuroma.
Nonmedullated nerve tissue,	Amyelinic Neuroma.
(*e*) Epithelial tissue tumours.	EPITHELIOMATA.
Squamous celled epithelium,	Squamous celled Adenoma. Squamous celled Papilloma.
Columnar celled epithelium,	Columnar celled Adenoma. Columnar celled Papilloma.
Spheroidal celled epithelium,	Spheroidal celled Adenoma.
Endothelium,	Angeioma. Endothelial Papilloma.
C. CYTOMATA.	
(*a*) Tumours of indifferent cells, (Blastocytes.)	BLASTOCYTOMATA.
(*b*) Tumours of connective tissue cells also of lymphoid tissue cells and muscle cells. (Mesocytes.)	SARCOMATA. (Mesocytomata.)
(1) Pure Sarcomata.	
Round cells,	Round celled Sarcoma.
Spindle cells,	Spindle celled Sarcoma.
Giant cells,	Giant celled Sarcoma.
(2) Compound Sarcomata.	
Fibrocytes,	Fibrosarcoma.
Chondrocytes,	Chondrosarcoma.
etc., etc.	etc., etc.
(*c*) Tumours of Epithelial cells.	CARCINOMATA. (Epicytomata.)
Squamous epicytes,	Squamous celled Carcinoma.
Columnar epicytes,	Columnar celled Carcinoma.
Spheroidal epicytes,	Spheroidal celled Carcinoma.
Endocytes,	Endothelial celled Carcinoma.
Syncytium,	Syncytial Carcinoma.

This classification has the great advantage that it enables us, if we are sure of what we see with the microscope, to place any given specimen in its proper place. In other words, the classification is based on facts which can be observed and not on any theories which may vary from time to time. Many attempts have been made by different authors to fashion a classification of tumours on an embryological basis but they all fail when put to the test. To my mind a classification based on embryology is unscientific and useless. A classification should always be based on facts and not on theories. We do not, for instance, classify common objects according to their ultimate derivation but according to their structure or the uses to which they are put. Who, for instance, would classify metal goods according to the mines from which the metallic ore was obtained? Yet this is exactly comparable to a classification of tumours on an embryological basis.

Another advantage of this scheme is the fact that the classes *Histiomata* and *Cytomata* correspond to the classes distinguished clinically as *simple* and *malignant* tumours. Each kind of tissue is represented by a tissue tumour and every kind of cell by a cell tumour.

NOMENCLATURE.

The different kinds of cells are named by adding the termination *-cytė* to a root expressing the nature of the cell according to the following scheme:—

A. Indifferent cells,			BLASTOCYTES
B. Epithelial cells,			EPICYTES.
C. Connective tissue cells,			
Fibrous tissue cells,	FIBROCYTES.	DESMOCYTES.	MESOCYTES.
Cartilage cells,	CHONDROCYTES.		
Bone cells,	OSTEOCYTES.		
Fat cells,	LIPOCYTES.		
etc.	etc.		
D. Bone marrow cells,	MYELOCYTES.		
Spleen pulp cells,	etc.		
etc.	etc.		
E. Free cells.			
Blood, Lymph, etc.	LEUCOCYTES.	ALETOCYTES.*	
	ERYTHROCYTES.		
	LYMPHOCYTES.		
F. Nerve cells,		NEUROCYTES.	
G. Muscle cells,		MYOCYTES.	

---

* Αλήτης Wanderer.

The histiomata are named by adding the termination *-oma* to a root expressing the nature of the tissue which is characteristic of the tumour. The adoption of this rule involves the abandonment of the term *epithelioma* as expressing one of the kinds of carcinoma. In this sense the term is wholely unnecessary and redundant and implies a distinction from other kinds of carcinoma which is not warranted by the pathology of the disease. For the same reason the term *endothelioma* is discarded in the malignant sense. Owing to the term *epithelioma* having been used as implying malignancy the epithelial histioma are called *papillomata* when the epithelium covers the surface, and *adenomata* when the epithelium forms tubes and spaces imbedded in a connective tissue stroma.

RELATION OF TUMOUR GROWTH TO OTHER PROCESSES.

There are numerous other diseases which bear a close resemblance to tumours in their general characters. They are characterised by the progressive nature of their course and by their apparently causeless origin.

In the breast we occasionally meet with a *progressive hypertrophy,* the whole breast enlarging without apparent cause and with no apparent limit to its growth. Similar cases are also met with in the thyroid (*Goitre*), in the prostate, etc.

Next we find cases in which one of the tissues of an organ, usually the connective-tissue, undergoes a progressive increase. This is seen in *progressive fibrosis* of nerves, in the prostate, and probably in some cases of cirrhosis of the liver.

The distinction between these conditions and histiomata is that in these conditions the whole organ is affected, the tissues retaining their typical relationship but being altered in amount, while a tumour is localised to one part of the organ. Progressive hypertrophy of, *e.g.*, a glandular organ corresponds to a diffuse adenoma while progressive fibrosis corresponds to a diffuse fibroma.

Again we find cases where the tissues undergo a *progressive atrophy,* as in the primary muscular atrophies, in progressive muscular atrophy, and in some other nervous diseases.

We also find cases of continued excess or defect of functional activity (*Hyperergasis, Hypo-ergasis*) as, for example, in diabetes insipidus, diabetes mellitus, exophthalmic goitre, and numerous nervous and mental diseases.

Lastly we find cases of *progressive metaplasia* as seen in Myositis ossificans and probably in the articular cartilages in Rheumatoid arthritis.

Other examples of one or other of these conditions are found in pernicious anæmia, erythrocythæmia, leucocythæmia, Hodgkin's disease, etc.

The processes underlying these diseases have always been a puzzle to pathologists and clinicians. They are closely allied to tumour growth. Like tumours these diseases arise without apparent cause or many different causes may appear to give rise to the same disease. Like tumours, also, their course is progressive and only slightly, if at all, influenced by treatment, and recovery is rare.

We must, then, consider tumour growth as a member of a group of processes which together may be called the *progressive processes* as opposed to the *reactive processes* which are seen in the inflammatory and adaptive changes.

DIFFERENCE BETWEEN SIMPLE AND MALIGNANT TUMOURS.

A most important point in connection with the classification of tumours is the following:—Is it possible by a histological examination to say definitely whether a given tumour is malignant or not? I have said that it is possible to allot any given tumour a place in the classification provided that we are sure of what we see with the microscope. In certain cases, however, it is not always easy to say exactly what the structure is at which we are looking. Endothelial cells resemble epithelial cells or sometimes connective tissue cells very closely and certain inflammatory conditions resemble certain tumours. Also certain carcinomata bear a close resemblance to certain adenomata. This difficulty can often be overcome by examining a number of sections from different parts and prepared by different methods.

In order to answer the above practical question satisfactorily it is necessary to decide on the question—What do we mean by malignancy in a tumour? The signs of malignancy are usually given as rapidity of growth, ulceration, infiltration, the production of metastases, and the presence of cachexia. To my mind the one necessary feature of malignancy is infiltration of the surrounding parts. The infiltration may be present in a very varying amount and may not be evident to the unaided eye but it can always be detected by the microscope if the right part of the tumour be examined. In other words, the distinction between the histiomata and the cytomata lies essentially in their mode of growth. If, for instance, we examine a chondroma we find everywhere typical cartilage surrounded by fibrous tissue (Fig. 3). In growing, it displaces the surrounding tissues without infiltrating them. In a chondrosarcoma, on the other hand, the nodules of cartilage, instead of being surrounded by fibrous tissue, are surrounded by sarcoma cells, and these cells are able to infiltrate the surrounding tissues, proliferating there and giving rise to the cartilage as a secondary product (Fig. 4).

Again, an adenoma presents itself as an encapsulated or pedunculated tumour and the epithelium of the tumour is everywhere surrounded by some of the stroma (Fig. 5), whereas a carcinoma, while it may resemble an adenoma in its older parts (Fig. 6), yet, in its actively growing parts, shows epithelial cells penetrating the interstices of the surrounding tissues, and any tubes or spaces in the carcinoma are of secondary origin (Figs. 7, 8). The tubes, once formed, are capable of continued growth in which they retain their tubular character. Hence, to be certain of the diagnosis in a carcinoma, especially in the columnar celled type, it is necessary to examine the spreading margin. Here it may be remarked that tumours do not always grow uniformly; one part may grow for a time actively and then cease while another part takes on active growth. Hence it may not be easy to find the particular part of the tumour, the study of which will give a decided answer to the question of the presence or absence of malignancy.

In order, then, that the pathological diagnosis may be of value, the pathologist must always be supplied with as much of the tumour as is available, including, if possible, some of the surrounding tissues, and he should be informed of the clinical features of the case. The use of the microscope affords the most certain, sometimes the only, means of diagnosing the nature of a tumour.

We can now reply to our question as follows :—Theoretically t is always possible by a histological examination to say whether a given tumour is malignant or not.

Practically it is possible in the great majority of cases to give a decided answer from a simple microscopical examination. In the few remaining cases it may be necessary to make an extended examination of different parts of the tumour before coming to a definite conclusion.

It is not always necessary to detect actual infiltration of the surrounding tissues to give a diagnosis of malignancy, as there are other factors which assist in the diagnosis, such as irregularity in the epithelium (in carcinoma), rapid and irregular proliferation as indicated by mitotic figures, etc., etc. It is often easier to say that a tumour is malignant than to give a decision on the less important question as to what type of cancer to refer it.

We see, then, that the cytomata differ from the histiomata in their mode of growth. In the former the spreading part is cellular, the formed tissues being of secondary formation, and in the latter the formed tissues grow as tissues from the first. In the former, the mode of growth is infiltrative, in the latter expansive. I shall have occasion to return to this point.

RUDIMENT OF ORIGIN.

In considering the life-history of a tumour we have, first, to consider what is the rudiment from which the tumour springs? What is the first beginning of a tumour?

Cohnheim endeavoured to explain the origin of tumours by supposing that they originated from rudiments which were sequestrated in embryonic life from the surrounding tissues, and took no further part in normal development. He supposed that these rudiments, retaining their embryonic characters, would, when the surrounding conditions were altered, as, for example, by irritation, show a greater power of proliferation, and he thus explained the great powers of growth exhibited by tumours. While this theory will not explain all tumours and is unnecessary in the great majority of tumours, it is applicable in certain cases.

Beard[3] attributes tumours to aberrant germ cells. According to this authority the fertilised ovum, in its initial proliferation, gives rise to a definite number of primitive germ cells, one of which gives rise to the embryo, while the others migrate into the body of the embryo and there become the primary germ cells of the sexual glands. Occasionally, instead of passing to the sex glands, these germ cells become deposited in other parts, where, in after life, they may give rise to tumours. There is not any evidence in support of this hypothesis, which is based on purely embryological grounds, and a strong objection to it is that it will not explain the close resemblance of a tumour to the tissues at its site of origin.

It was laid down in the definition of a tumour that the component parts of a tumour resemble structures normally found in the body. When we consider individual tumours we can expand this statement and say that the cells or tissues of which a tumour is composed resemble, in the great majority of cases, those present at the site in which the tumour originated. Seeing that this resemblance exists it is natural to suppose that the tumour arises, directly or indirectly, from the normal structures at the place of origin, and we can, I think, take it as a fact that this is the case in the great majority of tumours.

There are cases in which the tumours are composed of tissues not normally present at the site of origin, but even in these cases it is not always necessary to suppose an aberrant rudiment. We know, for instance, that under certain circumstances, it is possible for tissues to undergo metaplasia. The various kinds of connective tissue are interchangeable, as are also the various kinds of epithelium. We know, for example, that cartilage or bone may arise in connective-tissue: hence it is not necessary to suppose an aberrant rudiment of cartilage to explain the presence of a chondroma in positions in which no cartilage normally exists. Again, we know that columnar epithelium may undergo metaplasia into squamous; hence it is not necessary to suppose an aberrant rudiment to explain the occurrence of a squamous-celled carcinoma in the body of the uterus. I have seen a carcinoma of the cæcum which to a large extent consisted of typical squamous cells.

Ribbert[7] accepts Cohnheim's views for many tumours, but in addition states that the rudiments from which tumours originate may arise during post-natal life. In this case they consist, not of the normal tissues directly, but of some portions sequestrated from them by an inflammatory process or otherwise separated from their organic connections with the neighbouring parts, and these rudiments, being freed from the normal influences of the surrounding tissues, proliferate and so give rise to tumour growth. The first change in carcinoma, for instance, is a kind of inflammatory change in the connective tissue which proliferates and penetrating between the epithelials cells isolates them. These cells, being thus isolated and freed from the influences of the neighbouring cells, proliferate, and thus give rise to the carcinoma.

This idea, while undoubtedly true in many cases, is not, I think, necessary for all cases. It is a point very difficult to prove, and in some cases the connection between the tumour and the normal tissue is so close as to make it difficult to believe that the former has not developed directly from the latter.

We can say, then, that tumours arise from normal tissues directly or indirectly. They may also, in certain cases, arise from rudiments such as those postulated by Cohnheim. We may add that one tumour may arise from the tissues of another tumour.

The rudiment from which a tumour grows may be of any size. Some may start from a single cell, while others may originate from a whole organ, as in cases of diffuse carcinosis of the breast, stomach, or liver.

MODE OF GROWTH.

The next point to consider is: How does the tumour grow from the rudiment? This question will be considered more fully later on. We may here state that tumours grow by proliferation of their own cells, the surrounding tissues taking no part in the process. We must, however, limit this statement by saying that the area from which the tumour arises may increase in size, and, to this extent, the surrounding tissues may take some part in the growth, but it is only the tissues in direct continuity with the point of origin which act in this way. For instance, in a carcinoma of the breast the acini surrounding the tumour take no part in the growth, but are destroyed by it. Also in a carcinoma of the skin, while the area of origin may extend, yet the tumour itself grows by proliferation of its own cells, and, if it come in contact with the epidermis at a distance from the point of origin, the epidermis is destroyed by the advancing tumour and takes no part in the growth.

I have seen an angeioma of the liver which measured about 12 inches in diameter and reached as low as the iliac fossa hanging from the right lobe of the liver.[6] It was a cavernous angeioma and had been of rapid growth as far as could be determined from the history. Now throughout the tumour, even at its lowest margin, could be found, in the trabeculæ between the cavernous spaces, bile ducts and groups of hepatic cells (Fig. 9). The growth was too great to be due to a simple angeiectasis, *i.e.*, to a simple dilatation of pre-existing vessels, and, if the growth had proceeded by proliferation of the tumour cells alone we should not expect to find the hepatic remnants throughout it. Hence the most obvious explanation of the growth is that it started in a limited area, and, as growth proceeded, successive capillaries abutting on this area took on an active proliferation, and so the area of origin continued to extend among the liver cells, and, in consequence, the bile ducts and other hepatic remnants became elongated with the growth of the tumour.

Centripetal growth, then, does not occur except in connection with the area of origin, but the growth is centrifugal.

Centrifugal growth may be either central (expansive), occurring in the substance of the mass, or peripheral (infiltrating), occurring at the periphery. The former is the mode of growth characteristic of the histiomata, the latter of the cytomata. Peripheral growth, however, is usually accompanied by central or expansive growth, but in some cases the periphery may continue to grow while the centre has ceased growing or has become necrotic.

PART II.

CANCER

PART II. CANCER.

CYTOMATA.

Coming now to the cytomata with which we are more particularly concerned, you will observe that I have divided this class of tumours into three genera, *Blastocytomata, Sarcomata,* and *Carcinomata,* and I must define what I mean by these three kinds of cancer.

When a fertilised ovum divides, the cells which first result from its division are indifferent cells or *Blastocytes; i.e.,* they are cells which show no differences by which they can be distinguished from one another and, in the course of development, each may give rise to more than one kind of cell. As proliferation proceeds these cells become differentiated and, when this takes place, we can distinguish two types of cell:—*Epicytes,* or cells arranged in contact with each other forming a membrane which covers a surface or lines tubes and spaces (epithelial cells), and *Mesocytes,* which are cells separated from each other by intercellular substance or by the intervention of more or less definite fibrillar tissue (connective tissue cells, muscle and nerve cells). To these three types of cells correspond the three genera of the cytomata.

BLASTOCYTOMATA.* A blastocytoma is a tumour of which the essential cells are indifferent cells or *blastocytes*. The only means by which we can recognise these cells is the fact that they become differentiated into different types of cells, both epicytes and mesocytes. Hence the blastocytomata are mixed tumours containing epithelial and other elements both derived by differentiation from the fundamental indifferent cells. The best example of this kind of tumour is seen in the congenital mixed tumour of the kidney which is the commonest form of renal growth in children. If we study one of these tumours carefully we find that the most conspicuous elements are small round cells resembling sarcoma cells (Fig. 10). Embedded in these are numerous epithelial tubes and, here and there, can be found striated and smooth muscle fibres and the various forms of connective tissue. Now a careful examination reveals the fact that, in places, there is direct continuity between the round cells and the epithelial cells and it can be determined that the epicytes arise from the small round cells. The earliest stage in the process is seen as a circular grouping of some of the round cells (Fig. 11). Later on these cells assume the characters of epithelial cells and the centre of the group disappears leaving a lumen to the epithelial tube thus arising (Fig. 12), but still there is no sharp demarcation between the epicytes and the surrounding cells. The tube, once formed, continues to grow and develops a basement membrane. Hence, in many places, the epithelial tubes appear to be sharply marked off from the surrounding cells and to have no immediate connection with them. It is only in certain places that the connection can be traced. In other places a similar connection can be traced between the fundamental cells and the various other forms of tissue present (Fig. 13). In support of this view of these peculiar tumours is the fact that the epithelial tubes, while often showing branching, are independent of each other and lie free among the surrounding fundamental cells. This can be determined by means of serial sections.

---

* In the Erasmus Wilson lectures (10) I named these tumours BLASTOMATA. I have altered the name because objection has been taken to the use of the term Blastoma in this sense, this term having been previously used to include all tumours.

In the later history of these epithelial tubes it will sometimes be found that the wall becomes invaginated on one side and the epithelial cells become flattened so that a structure is produced which bears a striking resemblance to a glomerulus before any blood-vessels have penetrated into it (Fig. 14).

Another example of the blastocytomata is to be found in the mixed tumours of the parotid. In these tumours the fundamental cells are polygonal cells with branches which communicate with the processes of neighbouring cells. If we examine the preparation closely we find that, on the one hand, by approximation of the cells we get epithelial cells—the processes becoming prickles—and, on the other hand, by a separation of the fundamental cells by the intervention of a mucoid intercellular substance we obtain mucous tissue which in its turn may give rise to cartilage or fibrous tissue (Fig. 15).

Tumours with similar characters are found in the testis, the brain, and other situations.

It seems clear that in these tumours differentiation proceeds with proliferation and that the fundamental cells are indifferent cells, being capable of giving rise to various forms of differentiated cells. Hence the rudiment of origin of these tumours is a group of undifferentiated cells which retain their capacity for undergoing differentiation until the tumour develops. This is a rudiment in Cohnheim's sense. In the case of the blastocytoma of the kidney the rudiment would consist of the cells of the paraxial mesoblast which give rise to the intermediate cell mass from which the Wolffian body is developed and the development of the tubes in the tumour corresponds exactly to the development of the tubes in the Wolffian body and to the development of the glomeruli and convoluted tubes of the kidney which are of mesoblastic origin. The presence of striated muscle is accounted for by the close association of the intermediate cell mass with the mesoblastic somites from which the muscles are developed.

SARCOMATA. I have divided the sarcomata into two groups —the pure and the compound.

A pure sarcoma consists of cells and intercellular substance alone without the presence of any formed tissues, whereas the mixed forms contain one or more forms of definite connective-tissue in addition to the purely cellular portions, but these tissues are of secondary formation, being developed from the fundamental cells (Fig. 4). The pure sarcomata are subdivided according to the kind of cell present and the compound according to the different kinds of connective-tissue which are developed in them.

CARCINOMATA. The carcinomata, like the sarcomata, are subdivided according to the type of cell present, and we may, also, if we please, divide them into *pure carcinomata* in which no definite epithelium is present, the cells being arranged irregularly, and the *compound carcinomata—adenocarcinoma* or *papillocarcinoma*—in which definite epithelium is formed either lining spaces or covering outgrowths.

It will be observed that I include the malignant endothelial tumours among the carcinomata. I do this because endocytes bear a close resemblance to epicytes both in structure and arrangement, in fact endocytes are only a particular species of epicytes. We find that, both developmentally and under pathological conditions, epithelial cells may assume an endothelial character, and that endothelial cells may assume the columnar, cubical, or squamous type. Hence tumours arising from endothelium often bear a close resemblance to the columnar (Fig. 16), spheroidal (Fig. 17), and squamous-celled carcinomata (Fig. 18).

We see, then, that the three genera of the cytomata have similar characteristics. In all of them the fundamental elements are cells and in all of them tissues may arise as secondary formations. The pure blastocytoma, *i.e.*, a tumour composed entirely of blastocytes, probably exists, but we have no means of recognising a blastocyte apart from the differentiated cells which arise from it. Possibly the tumour which we know as the round-celled sarcoma is an example of a pure blastocytoma. The pure Carcinomata, *i.e.*, the carcinomata in which the epicytes are arranged irregularly without any tendency to form definite epithelium are seen, especially in the spheroidal-celled and in some sqamous-celled growths, and the pure sarcomata are the ordinary round-celled, spindle-celled and giant-celled sarcomata.

In the compound cytomata we recognise, in the blastocytomata, epithelium and connective-tissue, muscle, etc.; in the sarcomata various forms of connective-tissue; and in the carcinomata definite epithelium, either in the form of tubes (*adenocarcinoma*) or as papillary outgrowths (*papillocarcinoma*).

CANCER IN ANIMALS.*

Cancers and other tumours occur in all kinds of vertebrate animals and probably in some invertebrates (Figs. 19—24). They resemble in structure and characters the corresponding tumours in Man, and much valuable knowledge has been gained rom the experimental study of these tumours. This experimental study has been carried on especially in mice which are omparatively frequently affected by tumours and it has been ound possible to transplant from mouse to mouse carcinomata, arcomata, and chondromata so that the growth of the transplanted tumours can be followed from the earliest beginning. t is found that the tumour originating in a mouse as the result f inoculation with a small portion of a tumour from another mouse is the result of the proliferation of the cells of the portion f tumour introduced.

---

* For an account of cancer in animals see the various Scientific Reports of the Imperial Cancer Research Fund.

RUDIMENT OF ORIGIN.

With regard to the rudiment of origin of a cancer there is ittle to add to what we have already said in regard to tumours n general. Cancers may start from the normal tissues of the part, or from sequestrated rudiments, or again from other umours, and the rudiment may be single or multiple. In ertain cases a carcinoma may start from the epithelial cells of whole organ. These are, perhaps, more correctly described as ases of diffuse *carcinosis* of the organ. Such cases have been described in the breast and stomach and occasionally in the liver and other organs. Primary carcinoma of the serous membranes, particularly of the pleura, usually takes this form, appearing as a general thickening of the membrane involved. The same may be the case in sarcomata, *e.g.*, in diffuse sarcoma of the liver. These conditions bear the same relation to Carcinoma and Sarcoma that progressive fibrosis does to Fibroma.

The area of origin of a carcinoma may, in some cases, ncrease in size, successive neighbouring portions of epithelium aking on a cancerous growth, but, as a rule, the increase in ize, *e.g.*, of a cancerous ulcer, is due to continuous proliferation of the cells arising from the original rudiment, the surrounding epithelium being destroyed and taking no part in the tumour growth. Hence, from an examination of the tissues in the neighbourhood of a cancer, we cannot draw any conclusions as o the structures from which it has arisen.

As to the organs from which cancers arise we can say that he organs most frequently affected are those which are exposed to chronic irritation or traumatism or those which are liable to considerable variations in functional activity.

MODE OF GROWTH.

A cancer, like other tumours, grows by proliferation of its own cells. The predominant type of growth is, as we have seen, peripheral; that is to say, the tumour cells at the periphery proliferate and penetrate the surrounding tissues thus giving rise to the infiltration which is the characteristic feature of malignancy. In most cases, however, this peripheral growth is associated with central expansive growth, *i.e.*, it is not only the younger peripheral cells which proliferate but also the older cells in the mass of the tumour.

When this expansive growth occurs to a considerable extent the tumour may present an appearance of encapsulation due to the compression and consolidation of the surrounding tissues by the expanding mass. The encapsulation is, however, apparent only, and examination with the microscope will show that the peripheral cells are proliferating beyond the apparent capsule and infiltrating tissues. This apparent encapsulation is more frequent in Sarcoma than in Carcinoma while in Carcinoma more frequently than in Sarcoma we see the opposite condition, the peripheral cells proliferating and infiltrating while the central cells have lost their power of proliferation and are degenerating. Hence a carcinoma, as is seen sometimes in Rodent Ulcer, may continue to spread while the older parts become healed.

The peripheral mode of growth explains the destructive characters of cancers. The cells penetrating the surrounding tissues, find their way into all available spaces and proliferating there separate the constituents of the tissues and cause them to undergo atrophy *in situ* without displacing them (Fig. 25). Cancer cells can penetrate into and proliferate within fat cells (Fig. 26) and striated muscle fibres. Dense substances such as cartilage, in which there are no available channels, withstand the encroachments of a cancer for a long time. We can say, then, that cancers replace, rather than displace, the surrounding tissues.

MODES OF EXTENSION.

PERMEATION. Cancer spreads to the parts surrounding the rimary mass by a mode of extension known as *Permeation.* Ve have seen that the primary mass grows by the peripheral ells proliferating and infiltrating the surrounding tissues. ermeation is an extension of this process, the cells proliferating ontinuously along the lymphatic channels. In the case of xternal cancer this permeation of lymphatics takes place to a reater extent in the deep fascia than in the superjacent or ubjacent tissues while here and there processes of the growth row towards the surface and give rise to nodules of tumour aving no apparent connection with the primary mass. Thus re explained the cutaneous nodules sometimes seen in Carcinoma f the breast and also the infiltration of bones, muscles, etc.[4] his permeation is not confined to the lymphatic channels but nay take place along veins, tendon sheaths, nerve sheaths, ducts, tc., any natural channel forming a path by which the cancer an grow. While this proliferation takes place in continuity ve may find that the chain of cells is not continuous at any one noment. While the cells at the extremity of the chain are roliferating, some of the cells nearer the primary mass may ave disappeared and the channel been obliterated by cicatrisation. The direction of growth in permeation is not dependent on he direction of the current in the lymphatics, etc., but takes place o an approximately equal extent in all directions radiating from he primary mass. Secondary tumours may also be centres of ermeation.

Growth of the same kind, viz., proliferation in continuity, nay take place over long distances. We may find a tumour rowing along the large veins or along the whole length of the horacic duct.

METASTASIS. When the growth has penetrated into a ymphatic or vein it is easy to understand how individual cells r groups of cells may become detached and carried on by the ymph or blood-stream. Such detached cells, being carried to he smaller arteries or capillaries, become lodged as emboli and ommence to proliferate, thus giving rise to a new tumour with no direct communication with the primary growth. This phenomenon is known as *metastasis* and the growth as *metastatic growths*. The lymph glands, lungs and liver are naturally, owing to their position with regard to the circulation, the most frequent seats of metastatic growths. The entry of the cancer cells into the blood-vessels may be either directly, by penetrating the wall of a capillary or vein (Fig. 27), or indirectly by way of the lymphatics.

Certain organs, such as the alimentary canal and muscles, are very seldom the seat of metastatic growths. These organs may escape when there is widespread metastasis in other organs, and we must suppose that the tumour cells in these cases are carried to the alimentary canal and muscles as well as to the other organs. Apparently, however, the tumour cells in these situations do not meet with the conditions which are necessary for proliferation, and are therefore destroyed or remain latent.

Independent secondary growths may also arise from cancer cells being conveyed along the lumen of a canal, such as the alimentary canal or a duct, and settling in new situations. They may also follow upon the dissemination of cancer cells in the serous cavities.

Metastatic cancers do not usually show such an extreme power of infiltration as do the primary tumours. Often they are apparently sharply circumscribed and appear to be encapsulated. This is largely due to the fact that the expansive growth takes a larger share in the enlargement of the tumour than the peripheral, and hence the surrounding tissues are displaced rather than infiltrated.

Occasionally metastasis has been described in a tumour which does not show malignant characters histologically. Thus chondromata have been described as giving rise to metastases. In the great majority of these cases careful examination shows that the tumour is not a simple chondroma, but is a chondrosarcoma or a carcinoma with a cartilaginous stroma. Apparently, however, there are cases where this explanation will not apply and where the tumour is a pure chondroma and does not show infiltration. These cases are characterised by a distinct growth of the tumour along the veins, and, given the presence of a growing tumour in the lumen of a vein, it is not difficult to understand how portions can become detached and so orm emboli. The presence of the chondroma within the vein s due to the tumour having come in contact with the vein rom outside; ulceration of the vein, due to the pressure of the unyielding cartilage, then occurs, allowing the tumour to grow nto the lumen of the vein. These cases are, however, excessively rare.

MULTIPLE TUMOURS.

Multiple tumours are not as rare as they were once thought to be. The different tumours may be all of the same nature or may be of different types. They may be all malignant or all non-malignant or some may be malignant and others simple. Also the different tumours may arise in the same or in different organs and may even be combined into one growth.

I have, myself, seen several cases of multiple cancers. One of them was of particular interest. It was a case where a man had his eye removed for a melanotic sarcoma. Two years later he died with a large mass of growth in one of his ribs. Naturally this was supposed to be a metastatic growth but it proved to be a squamous-celled carcinoma apparently of endothelial origin, seeing that no primary tumour of this kind existed anywhere else in the body (Fig. 18). Besides this tumour he had innumerable spindle-celled sarcomata in the liver, each tumour being about the size of a pea and none of them pigmented.

Another interesting case was a columnar-celled carcinoma of the stomach in which the stroma was sarcomatous, being composed of small round cells. In this case there were several metastatic tumours all of which were sarcomatous. This appears to have been a case of primary carcinoma, in the stroma of which a sarcoma developed, the sarcoma becoming predominant.

Cases have also been described of mixed carcinoma and sarcoma in which the metastases have been some of one type and some of the other.

THE STROMA OF CARCINOMA.

A carcinoma is usually described as consisting of epithelial cells in a connective-tissue stroma, and the question arises whether both constituents are to be regarded as integral portions of the tumour. There can be no doubt that it is the epicytes which are the active constituents of the tumour, and the fibrous tissue is merely a supporting tissue derived from the connective-tissue of the organ in which the tumour is growing. The growing part of a carcinoma is entirely cellular, the epicytes infiltrating the surrounding structures (Figs. 25, 26), and the stroma is produced as a secondary formation from these structures by condensation and proliferation of the connective-tissue. This fact has been confirmed by the observation of experimental carcinomata in mice. In every case in which a carcinoma is transplanted from mouse to mouse the stroma degenerates, while the epicytes continue to live and proliferate. A new stroma is subsequently supplied by proliferation from the tissues of the animal into which the tumour has been transplanted.[2] There are cases, however, which it is difficult to explain on this view that the stroma is formed from the tissues in which the tumour is growing. Such are the rare cases of carcinoma with a cartilaginous stroma which gives rise to metastatic tumours of the same character. In some of these cases, at any rate, the tumour grows along the interior of a vein, and a portion may be bodily detached containing both epithelial cells and cartilage, both of which continue to grow when they become lodged in a new situation. We should have to assume in this case that the tumour was a combined tumour.

Whether this is so or not there is no doubt that the epicytes have an influence on the stroma and that the cells which give rise to a metastasis can so influence the tissues at the new site that they produce a stroma with the characteristics of the stroma of the original tumour. A carcinoma arising in any organ usually shows a stroma resembling the stroma normally present in that organ and the metastases show a similar stroma. This is especially seen in adrenal tumours. In the transplantation of mouse carcinoma the new formed stroma assumes the characters of the stroma of the original tumour.

I have mentioned that a sarcoma may arise in the stroma of a carcinoma and it has been observed in the experimental transplantation of carcinomata in mice that occasionally, during successive transplantations, a sarcomatous stroma appears and eventually supplants the carcinoma, the sarcoma itself being transplantable.

VASCULAR SUPPLY.

The blood-vessels in a cancer are derived from the capillaries of the part in which the cancer originates. As the tumour grows the capillaries enlarge and lengthen and new capillaries may be formed from them. At the same time the gradually enlarging tumour engulfs more and more of the surrounding capillaries and other vessels which then become available for its nutrition. The blood-vessels in a cancer retain the capillary structure to a large extent although they may become dilated into sinuses and they show little tendency to become fully developed into arteries and veins. Hence the vascular system of a cancer consists mainly of capillaries and wider sinus-like channels arranged in an irregular network which connect with the blood-vessels of the surrounding part at all points in the surface of the tumour.

In a carcinoma the blood vessels run in the stroma. In those cases of Carcinoma in which the stroma is present in very small quantities and in sarcomata where the vessels run along the cells, hæmorrhages are likely to occur owing to the imperfect structure of the blood-vessels. In endothelial cancers the tumour cells themselves may form new blood-vessels in the same way that the cells of a columnar-celled carcinoma may form epithelial tubes.

ULTIMATE DESTINATION.

Having considered the sources and modes of growth and extension of cancers we have next to consider what is the ultimate destination.

Unfortunately, the great majority of cancers continue to increase until death results. We have seen, however, that some cancers while spreading in one part heal in others. These cases are seen especially in Rodent Ulcer and atrophic Schirrus. Such tumours may continue to grow for years without producing any except local effects.

Some tumours, even sarcomata and carcinomata, disappear spontaneously. Several cases of undoubted cancer verified by histological examination have been described which have disappeared completely, either without any treatment, or with such treatment as could not produce such an effect of itself. We do not in the least know what are the conditions which lead to such a result. We do not know whether there is a spontaneous cessation of proliferation or whether there is an enhanced effort on the part of the organism to stop the encroachment of the tumour. From the rarity of these cases it would seem to be an idiosyncrasy on the part of the individual tumour or patient and we have no means at present of reproducing the conditions in other tumours or other patients.

Some light, however, has been recently thrown on this question by the experimental investigation of mouse cancer. It is not uncommon for a cancer in a mouse, after growing for some time, to cease growing and to degenerate, ultimately disappearing completely. Such mice are found to be immune to subsequent inoculation. This immunity is specific in the sense that a carcinoma does not confer immunity towards a sarcoma and *vice versa,* and it is specific in a stricter sense. It is found that a given carcinoma confers a much higher degree of immunity towards subsequent inoculation of the same tumour than towards a carcinoma of different origin.

DEGENERATIVE PROCESSES.

Apart from disappearance, cancers may undergo various retrogressive processes, such as fatty and colloid degeneration and calcification. These changes in cancers are not always to be regarded as strictly degenerative changes. We find that the alterations often correspond to the changes which take place normally in the cells from which the cancer has arisen, and the changes are thus more allied to secretory than to degenerative processes. Fatty changes are especially common in breast cancers; the cells of a carcinoma of the thyroid secrete colloid; columnar-celled carcinomata of the stomach and intestines secrete mucin; those of adrenal carcinomata contain fat and cholesterin. Also sarcomata of bones are prone to ossify or calcify.

Cancers can also exhibit inflammatory changes.

METABOLISM.

We know little about metabolism in Cancer. Seeing that the cancer cell is derived from the cells of the organism we should naturally expect that, making allowance for the rapid proliferation, the metabolism would be similar to that of the normal tissues of origin. I have mentioned that carcinoma cells often secrete the same substances as the normal cells of the organ in which the carcinoma arises. Besides the examples given above I may mention that trypsin has been found in carcinomata of the pancreas and pepsin in carcinomata of the stomach.

Many carcinomata contain a large amount of glycogen, and some have maintained that the amount of glycogen is proportional to the degree of malignancy.* This is, however, not necessarily the case. While it is common in cancers, it is also found in some simple tumours, being always present in chondromata. The presence or absence of glycogen appears to depend more on the origin of the tumour than on the rapidity of its growth.

There do not seem to be any definite changes in the urine, temperature, pulse etc., due to cancer *per se*.

In studying questions of metabolism it is difficult to separate the effects of the cancer *per se* from those due to the disturbance of the organ affected.

CLINICAL FEATURES.

The clinical features of cancer can be explained by the pathology of the disease. The fixation of the tumour to the surrounding parts is explained by the infiltrative mode of growth. The appearance of cutaneous nodules in the neighbourhood of an external cancer is attributable to the permeation of the lymphatics. The extension to the lymphatic glands is attributable to permeation and embolism, and visceral deposits depend on metastasis by the blood-stream. Degeneration and softening depend on the imperfect blood supply, and cachexia on the drain of nutrient material combined with the absorption of the products of disintegration of the tumour cells, and of bacterial products in the case of ulcerated cancers. There does not seem to be a specific cancer toxin.

Recurrence after removal is due to imperfect removal, and is not characteristic of cancer alone. Simple tumours recur if the removal is incomplete. In cancer, however, since the tumour is not encapsulated, it is impossible to define its bounds, and hence outlying cancer cells are very liable to be left behind to give rise to a recurrent growth. Also, since cancer cells are transplantable, infection of the wound is liable to occur at the operation. When recurrence occurs at a distance from the original operation it is not necessarily to be attributed to imperfect operation. We have seen that multiple tumours are not so very uncommon; hence when recurrence takes place at a distance it is quite likely that the new tumour is independent altogether of the original one. This is especially likely to be the case when the new tumour arises in a part which is a common site of a primary tumour. When, for instance, after removal of a breast, a cancer appears in the other breast, it is possible, and even probable, that this second tumour is a new formation altogether and independent of the former one. The breast is a rare situation for a secondary tumour, while it is a common site of primary growths. In some of these cases of recurrence after operation the new tumour shows a different histological structure from the original one, and hence must be regarded as independent.

PART III.

CAUSATION

PART III. CAUSATION.

CAUSAL FACTORS.

HAVING considered the rudiment of origin and the mode of growth by which the cancer or other tumour grows from the rudiment, we have next to consider why tumour growth occurs or, in other words, what is the cause of Cancer?

The causal factors which give rise to any disease can be divided into the *essential* causal factors, or those factors without which the disease in question could not arise, and the *adjuvant* causal factors, or those factors which assist in the causation of the disease without being, themselves, individually necessary. Causal factors may also be divided into the *extrinsic* and *intrinsic* factors.

We have, then, two classes of diseases:—(1) Those in which the essential causal factors are extrinsic; (2) those in which the essential causal factors are intrinsic. A disease may be specific in two ways:—(*a*) It may be characterised by a specific extrinsic causal agent; (*b*) it may be characterised by specific lesions. If we consider a disease as characterised by a specific extrinsic causal agent we find that it does not show constant lesions. If, on the other hand, the disease is characterised by specific lesions it is not characterised by a specific extrinsic causal agent.[14] In other words, any extrinsic causal agent will, under differing circumstances, give rise to different lesions, and any given lesion may be due to many different extrinsic causal agents. For instance, pneumococcus infection may be characterised by fibrinous, serous, or purulent inflammations of various parts of the body, and, on the other hand, fibrinous pneumonia may be due to other organisms than the pneumococcus.

EXTRINSIC CAUSAL FACTORS.

It has long been known that extrinsic factors play a part in the causation of Cancer. Cancers may arise in chronic ulcers, in patches of lupus, or in scars derived from these. They may also arise in connection with injury as in the case of Sarcoma arising in connection with a fractured bone, or in connection with chronic irritation from a clay pipe, a jagged tooth, gall stones, irritating food, soot, etc.

The chief theory which associates Cancer with extrinsic factors is that which attributes it to the action of specific parasites.*

While at first sight, Cancer has a close resemblance to such a parasitic disease as Tuberculosis, yet, on closer examination, we find that this resemblance is no ground for concluding that Cancer is due to a parasite. Tubercle bacilli introduced into the tissues give rise to an inflammatory reaction with the production of fibrous tissue. Similarly cancer cells give rise to an inflammatory reaction with the production of fibrous tissue. Hence the cancer cell is analogous to the tubercle bacillus and we cannot therefore argue that the proliferation of cancer cells is due to a parasite. If we attempt to compare the tubercle bacillus with the supposed cancer parasite the analogy, and the argument derived from it, fail completely. In fact Cancer may be considered as an infective disease only in the sense that the cancer cell is itself the infecting agent.

A serious objection to the parasitic theory is found in the impossibility of explaining on this ground the congenital cancers and the non-malignant tumours and yet there is no reason to suppose that the causation of Cancer is essentially different from the causation of other tumours.

---

* For a criticism of this theory see appendix.

Parasites may give rise to tumours but they only do so indirectly in a similar way to other irritants and not in any specific manner. Examples are seen in the tumours arising in connection with Lupus and Bilharziosis.

The extrinsic causal factors are thus adjuvant and not essential factors.

INTRINSIC CAUSAL FACTORS.

We have now to consider the intrinsic factors in the causation of Cancer and other tumours. This is a more difficult subject than the extrinsic factors since we know very little about the intrinsic factors underlying normal conditions of growth, etc. To understand this question we must go back to the first principles of biology.

CELL PROLIFERATION.

There have been many discussions as to whether the phenomena of life are to be explained by the known forms of energy or whether a special *vital principle* is necessary for their explanation, and physiologists have always been divided on this point. Personally, I believe that there is a vital energy which is as distinct a form of energy as electricity. This vital energy manifests itself especially in assimilation and proliferation. While it can only be derived from pre-existing vital energy, it is maintained and increased by the transformation of other forms of energy, and, in its turn, it can manifest itself as mechanical energy, heat, light, electricity, or chemical action.

The chief characteristic of living organisms is proliferation, and this power of proliferation is inherent in all living beings. All living organisms continue to multiply except so far as they are prevented by antagonistic forces.

In Nature the restraints are of varying character. The higher organisms are restricted to a large extent by the necessity for sexual congress. All organisms are subject to the restraint arising from the struggle for existence, especially in the difficulty of obtaining the necessary nutriment and in resisting the attacks of their enemies, and, in the course of evolution, it has come about that those organisms which are most liable to destruction have developed the greatest proliferative capacity, as in the case of bacteria, animal parasites, fish, etc.

Now what is true of living organisms in general is equally true of the constituent parts of the organism, that is, of the cells which make up the structure of the organism.

We find that all the cells of the organism are capable of proliferation in a varying degree. All proliferate during the development and growth of the organism. After a certain time, which differs in different organs, we find that some cells, such as neurocytes, have practically lost the power of proliferation; others, such as myocytes and desmocytes, retain the power of proliferation under certain conditions, while some epicytes and the mother cells of leucocytes and erythrocytes continue to proliferate throughout life to make good the loss due to various causes. Ova and spermatozoa lose the power of proliferation on attaining maturity, to regain it only by subsequent fertilisation.

Now the sites of origin of tumours correspond accurately to these facts. We find that neurocytes which have lost the power of normal proliferation only give rise to tumours with the greatest rarity; striated myocytes also extremely rarely give rise to tumours, while tumours arise from smooth myocytes much more frequently, and especially in parts in which, like the uterus, they show a physiological multiplication at intervals throughout life. The cells which most frequently give rise to tumours are the epicytes, and here again the most frequent sites are those where proliferation occurs regularly, or at intervals, throughout life such as the skin, alimentary canal, breast, and female organs of generation, while those epicytes which are normally more quiescent, as in the liver, kidneys, sweat glands, etc., only rarely give rise to tumour formation.

Proliferation, then, is an inherent property of the cell, as it is of the living organism, and those cells which have the greatest proliferative capacity are the most frequent sources of tumour growth. These facts tell strongly against the theory of embryonic rudiments as an explanation of tumour growth.

INFLUENCES WHICH LIMIT PROLIFERATION.

We have next to consider why tumour growth does not occur as a general rule, or, in other words, what are the influences which limit proliferation under normal conditions? This is another way of stating the question, What are the influences which determine the form and size of any living being?

When the ovum undergoes segmentation the cells at first formed resemble one another. Soon, however, differentiation takes place and the cells assume characters which distinguish them from each other.

The different characters which the cells then assume appear to depend to a large extent on their position in the embryo, or, in other words, on their position in relation to the other cells of the embryo. Since the fertilised ovum contains all the potentialities of the organism, differentiation would appear to consist in the progressive loss or latency of certain characters and a consequent predominance of the characters which remain. Differentiation is found to a certain extent in some of the colony-forming protozoa in which some of the individuals have motile functions, others nutritive, others generative.

The cells, in spite of differentiation, retain their power of proliferation which is inherent in them, but different kinds of cells retain this power in very varying degrees.

CO-ORDINATING MECHANISM OF THE ORGANISM.

Seeing, then, that this differentiation and proliferation occur in the different tissues, we have to ask ourselves, What is the co-ordinating mechanism of the organism? What are the influences which so regulate differentiation and proliferation that a perfect condition of equilibrium is maintained? Even in those cells which proliferate continuously there is a condition of equilibrium, because the proliferation is just sufficient to balance the corresponding destruction. In the skin and mucous membrane the epicytes proliferate only to such an extent as to replace the continual waste of cells which are cast off at the surface, and in the case of leucocytes and erythrocytes the production of new cells is balanced by the destruction which takes place in the spleen and elsewhere. In those tissues, the cells of which do not show continuous proliferation, we find that the destruction of some cells is followed by regenerative proliferation of the cells which remain, which ceases when the defect is made good.

This question of the co-ordinating mechanism of the organism is the fundamental question for the pathology of tumours. If we knew what are the influences at work in producing the co-ordination, then we should be very near solving the problem of the causation of tumours, since tumours arise from a defect in co-ordination. Unfortunately we do not know what these influences are.

The form of any organism, which depends on differentiation and proliferation, is an inherent property of the species and is transmitted by heredity. Hence the co-ordinating influences are a property of the germ plasm and are transmitted equally by the ovum and spermatozoon. The influences within the organism are apparently intercellular, each cell exerting certain influences upon its neighbours. That such intercellular influences exist is shown by the co-ordination of the movements of the cilia in ciliated epithelium. This ciliary movement remains regular and co-ordinated in a detached piece of ciliated epithelium for a long time. As to the paths by which these influences pass from cell to cell, it would appear to be by direct continuity between the cytoplasm of adjoining cells, each cell being in protoplasmic continuity with its neighbours. This has not been entirely proved, but we can see the connections between the prickle cells of the epidermis, and they have been demonstrated in other kinds of epithelium, in smooth muscle, in bone, in fibrous tissue, and, according to some, they are present in cartilage.

In fact we have come to take too strict a view of cellular pathology. We are accustomed to explain all pathological changes by cellular activities without considering the organism of which the cells are the constituent parts. A living organism, however, is not a mere aggregation of cells, it is a distinct entity in itself—an individual—and the cellular activities are everywhere subordinated to the control of the whole organism.

ıe multicellular organism can be regarded as being composed cells, differing in structure according to the function they .ve to perform, which everywhere retain intercellular proto-asmic connections, and it is through these connections that the -ordinating mechanism of the organism acts. Development d growth are to be considered as primary properties of the ganism as a whole, the proliferation of the cells being bsidiary. That is to say, the organism does not grow because e cells proliferate, but the cells proliferate because the ganism grows.

Under normal conditions cells and tissues separated from eir organic connections can form secondary connections with e neighbouring parts. Thus, when leucocytes penetrate tween prickle cells, the intercellular bridges which they break their progress are reunited behind them. Also in skin graft-g and the reunion of wounds connections are formed with e neighbouring parts, and by means of these secondary con-ctions the organism is again able to exert its co-ordinating fluence and so to limit the cellular proliferation to the amount cessary to repair the deficiency.

The intercellular influences, besides regulating proliferation, so play a part in controlling functional activity.

While this co-ordinating influence is thus a property of e organism as a whole, it does not appear to be centred in y particular situation. It is also shown during the survival detached portions, especially in the lower organisms. Hence portion of one of the lower organisms can reproduce the ıole, a heart removed from the body will continue to beat ythmically, a fragment of ciliated epithelium still shows -ordination of the ciliary movement.

THE RELATION OF THE TUMOUR TO THE ORGANISM.

If a portion of tissue, or group of tissues, be separated from organic connection with the neighbouring parts, while retain-g its position in the organism, and be unable to form secondary nnections, it will no longer be subject to the control of the ganism. Therefore, if it begin to grow, it will grow as an dependent individual, dependent on the organism only as far as its blood supply is concerned, thus forming a tissue tumour. Growing as an individual it will displace the surrounding tissues as it increases in size but, having no reproductive mechanism, it will not give rise to secondary tumours. If, however, a piece of the tumour be detached, as in the rupture of a papilliferous ovarian cyst or in a piece left behind in incomplete removal, such a detached portion will itself grow into a tumour resembling the old one provided that the vascular connections remain sufficient. While the tissues of which a simple tumour is composed are removed from the influences of the organism, they retain their connections *inter se* so that the whole tumour grows as an individual.

A simple tumour can thus be regarded as an asexually produced descendant which lives parasitically in the parent organism.

In a cancer the case is different. Here it is a cell which is freed from the co-ordinating influences of the organism. The cell thus proliferates, giving rise to new cells which are also independent. These cells penetrate into the spaces of the surrounding tissues, and wherever they settle they give rise to new groups of cells. A cancer, then, cannot be regarded as an individual; it is a colony of cells and the cells themselves are individuals.

The objection may, perhaps, be raised in this connection that the blood corpuscles of the normal body are independent cells and yet do not show the proliferative capacity which we find in cancer cells. The blood cells, however, are in no sense tissue cells. They are cells which, when they have become free, are destined for a limited existence only and do not proliferate under normal circumstances, except to a slight degree by amitosis.

A simple tumour, then, is to be regarded as an asexual multicellular organism while a cancer is comparable to a colony of unicellular organisms, and, just as unicellular organisms can form among themselves secondary associations with definite characters, as is seen in bacteria and the colony-forming protozoa and diatoms, so the cancer cells can form secondary associations *inter se* with the formation of distinct tissues. We find that hose cancers in which these secondary associations are formed nost readily are the least malignant and *vice versa*, those cancers n which the cells remain independent of each other are the nost malignant. For instance, among carcinomata the cells of the columnar-celled type most readily form tissues (epithelial tubes) and this is the least malignant species. On the other hand the cells of a spheroidal-celled carcinoma show very little tendency to form associations *inter se* and we find that the spheroidal-celled carcinoma is the most malignant form.

Another character in which cancer cells resemble protozoa s the limitless capacity for proliferation which they exhibit. Jensen's mouse cancer has now been growing for over seven years, that is, for a far longer period than the duration of life in a mouse, and it shows no sign of diminishing in vigour. It has been transplanted into innumerable generations of animals, and descendants of the original tumour are now to be found in laboratories throughout the world. At each successive transplantation the epithelial cells of the implanted tumour continue to proliferate and so give rise to the new tumour, so that the cancer cells show the same limitless proliferative capacity that is found in unicellular organisms.

Whether or not cancer cells show a still closer resemblance to unicellular organisms by occasionally manifesting a sexual process in the course of reproduction is not certainly known, but various observations on the characters of the mitoses in cancers tend to show that such is the case although some of the changes seem open to doubt. Appearances are sometimes found which suggest that cancer cells sometimes conjugate.[2]

Cancer cells show a further resemblance to unicellular organisms in that they will withstand the effects of cold. The cells of a mouse cancer can be kept at a temperature of 0°C. for several days or weeks,[2] and will even withstand the temperature of liquid air for some time and still retain the power of giving rise to a tumour when introduced into a new mouse,[3] [1] whereas if kept outside the living body at body temperature they quickly die.

A malignant tumour, then, is not a distinct entity in itself, it is a colony of cells, and Cancer can truly be regarded as a process of infection by cancer cells.

The growth of a cancer thus differs from the growth of the organism. In the organism the proliferation of the cells is secondary to the growth of the whole cell-complex, while in a cancer the growth of the whole is the result of the proliferation of the cell units.

If Cancer be regarded in this light all the phenomena of malignant disease are easily understood.

THE NATURE OF THE INTERCELLULAR INFLUENCES.

I do not intend to discuss here the phenomena of mitosis but I may remind you that cell division is initiated and effected by the centrosome with its associated amphiaster. These structures are formed of a substance known as *archoplasm*. It is not certainly known whether the archoplasm is a permanent constituent of the cell or whether it is modified cytoplasm, but there are reasons for believing that it is permanent and that the centrosome and archoplasm are handed down from cell to cell in the same way as the nucleus and cytoplasm. The centrosome appears to control not only the movements which result in cell division, but also amœboid and ciliary movements. Thus, while the nucleus and cytoplasm are concerned with nutritive and metabolic changes, the centrosome and archoplasm are concerned in the movements and division of the cell. It may be, then, that the intercellular connections are composed of archoplasm.

As to the nature of the influences which pass from cell to cell we know nothing. We can only say that they are an inherent property of the species and are handed down from generation to generation by heredity. They are present and active from the first commencement of division of the fertilised ovum, and hence cannot depend altogether on the presence of substances which are elaborated in organs which are developed at a later period, unless a store of such substance is laid up in the ovum sufficient to control development and growth in the early stages.

While it may be that some chemical substance, by reason of its chemical or physical properties, may play a part in controlling ell proliferation, it is difficult to imagine how such a substance ould control the form and arrangement assumed by the mass f cells resulting from the proliferation. It is inconceivable hat the form of any species of organism can be dependent on ne presence of any chemical substance. The controlling inuences would seem to be of a more subtle nature. It is in onnection with questions like this that we seem bound to admit ne existence of a special form of energy which is peculiar to ving organisms.

Chemical substances may, however, play some part in the ontrol of growth, just as they take part in co-ordinating unction in certain parts of the body. Indeed it is known that ne secretions of the ductless glands have some influence on rowth.

I have lately been investigating the nature and significance f certain crystals which occur in tumours. If a section of a ancer, which has not been subjected to alcohol, be examined y polarised light, crystals are found in nearly every case. ome of these crystals are fats or fatty acids and others are holesterin. In addition there are minute needle-shaped or rismatic crystals occurring singly or in clusters, either in or mong the cells. These crystals are distinguished by their eculiar behaviour on melting and subsequent cooling. Under ese conditions they do not again solidify but assume a cryalline fluid condition appearing as doubly refracting globules ig. 28). These crystals are found especially in and among e healthy proliferating cells of the tumour while the crystals fat and cholesterin are found especially in the degenerated eas. Degenerated areas are, however, often free from crystals any sort while the proliferating areas may contain numbers them.[12]

Having found these crystals in a number of cancers and in me other conditions, I made an extended investigation into eir nature.[11] It can be shown that they contain cholesterin nd, in some cases, fatty acids. I therefore experimented ith pure cholesterin and other chemical substances in der to find out under what conditions the peculiar doubly fracting globules could be obtained. I found that if cholesterin was melted with one of the higher fatty acids or certain alcohols and other substances, the globules appeared during the cooling of the preparation, and that the mixture subsequently solidified in the form of crystals closely resembling the crystals found in the sections.

It is fairly certain that these crystals in the tissues consist of cholesterin in loose combination with other substances. What the associated substance is cannot always be detected. Often it is a fatty acid as shown by the staining reaction of the globules resulting from melting the crystals; at other times they do not show the fatty acid reaction. It is probable that lecithin is one, perhaps the commonest of these associated substances, but we have no reliable means of detecting lecithin with the microscope. When these cholesterin mixtures are in the crystalline fluid condition they display certain peculiar physical properties. They can form an emulsion with water and it appears that cholesterin exists normally in the body as a colloidal solution or fine emulsion in association with lecithin or fatty acids, the cholesterin and the associated substance helping to keep each other mutually in solution in the body fluids. It is certain that cholesterin plays an important part in cell life, and the presence of these crystals in cancers, apart from any evident degeneration, suggests that cholesterin may be associated in some way or other with the regulation of cell proliferation.

As bearing on this point I may mention that spermatozoa are very rich in cholesterin, a fact which points to its being of importance to the developing embryo.

Another interesting point in this connection is the fact that gall stones are 2½ times as common in patients with carcinoma as in patients of similar ages suffering from other diseases.

The origin of Cholesterin is not certainly known but, from my observations, it would seem probable that it is secreted by the cortex of the adrenal body. I have examined a large number of adrenals and I find in every case crystals, resembling the crystals found in cancers, consisting of Cholesterin in association with other substances (Figs. 29, 30, 31). Usually these crystals occur in great masses filling the whole cortex and giving rise to the condition often called fatty degeneration of the drenal. It is, however in no sense a degeneration. It appears o be a normal condition and it seems as though the function of he adrenal cortex is to secrete Cholesterin and to pass it into he blood.[13]

Now we find that maldevelopment of the adrenals is usually ssociated with maldevelopment of other parts, especially of the rain, or with infantilism. On the other hand, hypertrophy nd tumours of the adrenal cortex in children are associated with sexual precocity or with gigantism. Hence it would appear that the secretion of the adrenal cortex has some share in the mechanism of the control of growth and development. These observations, then, suggest that Cholesterin is elaborated in the adrenal cortex and that it plays some part in the regulation of cell proliferation.

I may mention that, some years ago, Mr. Holden Webb of Melbourne, suggested that Cancer was due to the crystallisation of Cholesterin from the living cells and he proposed a method of treating Cancer by hypodermic injections of soap solution, the idea being that the soap would help to keep the cholesterin in solution. Mr. Webb's suggestion appears to have been pure hypothesis as he did not detail any evidence on which it was founded.[9]

PHYSIOLOGICAL EQUILIBRIUM.

We have next to determine why an extrinsic causal factor, which normally gives rise to inflammatory or compensatory changes, occasionally gives rise to tumour formation.

An epithelial cell is acted upon by the influences of the contiguous epithelial cells and also by the influence of the subjacent connective tissue. Normally these influences are in equilibrium with the intercellular proliferative forces. At the free surface, there being no antagonistic influences, the activity of the cell is manifested by proliferation, as in the skin and mucous membranes, or by secretion as in glands, or by absorption. Similarly a connective tissue cell is subject to the influences of the neighbouring cells which are in equilibrium with the proliferative forces within the cell.

Now the changes that occur when the equilibrium is disturbed will differ according as the original condition of equilibrium was *stable* or *unstable*. If the condition was stable any disturbance, by removing the influences, will be followed by proliferation of the cell which will continue until the influences of the neighbouring cells are again restored when the proliferation will cease. On the other hand, if the original condition of equilibrium was unstable, the intercellular influences will not be restored and proliferation will continue. Take, for example, an epithelial cell, the proliferative forces of which are in equilibrium with the restraining influences of the neighbouring cells. If we disturb the equilibrium by removing some of the neighbouring epithelial cells, the proliferative capacity is set free in the direction of the removed influence. The cell therefore commences to proliferate and, if the original condition of equilibrium was stable, proliferation will continue until, by the replacement of the cells which had been removed, the intercellular influences are again restored when proliferation will cease. This sequence of events we see in the regeneration of epithelium which follows an abrasion or a wound. If, however, the original condition of equilibrium was unstable, the intercellular influences will not be restored. Hence proliferation will continue indefinitely thus giving rise to a papilloma or an adenoma. If, instead of removing the influence of the neighbouring epithelial cells, we disturb the equilibrium by removing the influence of the subjacent connective tissue, then, if the original condition of equilibrium was stable, the cell proliferates for a time and then ceases as the restraining influences are restored. This is seen in the traumatic epithelial pearl produced by the displacement of epithelial cells into the subjacent connective tissue. If, on the other hand, the original condition of equilibrium was unstable, the restraining influences will not be restored and proliferation will continue in the connective tissue thus giving rise to a carcinoma.

Similarly in regard to the connective tissue cell, disturbance of the equilibrium, if stable, leads to regeneration and repair, but if unstable the result is a histioma if the relation between the cells and intercellular substance continue normal, or a sarcoma if the cells proliferate without producing the intercellular substance.

Similar conditions occur with regard to functional activity. The functional equilibrium, just as the structural, may be stable or unstable and the result of disturbance is accordingly compensatory or progressive Hyperergasis.

We can say, then, that a tumour arises from a disturbance of a position of unstable equilibrium between the proliferative forces within the cell and the antagonistic influences of the neighbouring cells. The characteristic feature of an unstable condition of equilibrium is that it needs only a very slight force to upset it, hence the same extrinsic factor which, under normal conditions, only produces a temporary disturbance, will, if the equilibrium be unstable, give rise to tumour formation. An injury to a bone usually gives rise to a fracture with subsequent healing or to a periosteitis with subsequent recovery, but a similar injury may, in some cases, be the starting point of a sarcoma of other tumour.

The instability of equilibrium is probably a local condition as a rule but it is possible that it may occur as a general diathetic condition.

While the instability of equilibrium will explain the origination of a tumour it will not explain its extension to distant parts and its continued growth in metastatic nodules and in transplanted tumours. We have seen that cancer cells must be regarded as unicellular organisms which, while they may form new associations among themselves, have lost the power of forming secondary connections with normal tissues through which restraining influences could pass. Moreover it is possible that the cells, by their escape from the control of the organism, have acquired an increased capacity for proliferation—a habit of growth—which continues in the sites of metastatic nodules. The cancer cells therefore continue to proliferate as long as they are supplied with the necessary nutriment and the reaction of the surrounding tissues is of the same kind as the reaction caused by parasites or foreign bodies. The growth in metastatic nodules is usually more circumscribed and less infiltrative than at the primary site where the condition of unstable equilibrium exists.

The immediate essential causal factor of tumour formation is thus an *unstable condition of equilibrium* between the component parts of the body, and the next question we have to consider is: What are the causes of this instability?

THE INFLUENCE OF INFLAMMATORY CONDITIONS.

It has long been known that chronic inflammatory conditions may lead to Cancer. A chronic ulcer of the leg may continue for years without appreciable alteration, but the continued irritation may at length so alter the condition of the tissues that the equilibrium becomes unstable, when the result is tumour formation. Similarly Carcinoma may start in a chronic gastric ulcer, or in connection with the chronic inflammatory changes produced by Lupus or Bilharziosis, and Sarcoma may arise from the callus of a healing fracture. Also Leucoplakia of the tongue is said to be a frequent precursor of Carcinoma. The irritants in these cases do not act directly on the epithelial cells, but they so alter the connective tissue that the equilibrium between it and the epithelium becomes unstable so that tumour formation follows. In Bilharzial Carcinoma, as Dr. Harris pointed out, the ova, which are the irritating factors, are not situated in the growth itself but in the neighbouring connective tissue.[5] (Figs. 32, 33.)

Other examples of chronic irritation producing cancers are seen in the influence of clay pipes, jagged teeth, etc., on the occurrence of cancer of the mouth.

During the proliferative changes which accompany these inflammatory conditions it is very likely that certain tissue cells will become separated from their normal connections. Such isolated cells may form the rudiments from which a tumour may arise, either during the progress of the inflammation, or, after it has healed, from the scar.

THE INFLUENCE OF HEREDITY.

It is impossible to obtain reliable statistics as to the part played by heredity in cancer because it is only in rare instances hat we can obtain a reliable family history. There is no doubt, owever, that cancer does occasionally run in families and this s what we should expect.

In many cases, however, the tendency to cancer in an individual is not so much due to heredity as to a want of heredity—t may arise in the individual as a spontaneous variation or nutation. It is these spontaneously arising cases that we should xpect to be handed on to the descendants by heredity since it s an accepted fact that variations tend to be inherited. On the ther hand, cases of acquired cancer, such as those arising in onnection with chronic irritation, we should not expect to be ransmitted to descendants as it seems clear that acquired haracters are not inherited.

THE INFLUENCE OF CIVILISATION.

There can be little doubt that the various influences grouped ogether under the title of civilisation play a part in producing tendency to Cancer. It is difficult to obtain statistics with egard to Cancer in uncivilised races and wild animals but it eems certain that, while Cancer is not unknown under these onditions, it is much more rare than among civilised races and omesticated animals. It is probable that no one factor in ivilisation is responsible but that it is the condition as a whole hat is at fault. Unnatural and excessive food, unhealthy surroundings, indoor and sedentary occupations, and the mental nxiety and worry which are inseparable from civilised life robably all take a share in producing a wear and tear of life hat is conducive to the formation of an unstable condition of quilibrium.

THE INFLUENCE OF SEX, AGE, AND SITE.

When we come to study questions regarding the incidence of Cancer, we are met with the difficulty of obtaining reliable statistics from which we can deduce conclusions. To be of any value the statistics must deal with a sufficient number of cases to eliminate accidental fallacies and each case must be verified microscopically by a competent histologist. Even if this precaution be taken a further difficulty arises in the fact that pathologists do not always agree as to the position of certain tumours. Endothelial tumours will by some be placed among the sarcomata and by others among the carcinomata, while others will allot them a class by themselves. Also some will include gliomata and myxomata with the sarcomata. The majority of the tumours which I have called Blastocytomata are included by others among the sarcomata although the fact that epithelium develops in them distinguishes them from the true sarcomata. Many place the mixed tumours of the parotid among the endotheliomata, without any good reason as far as I can see.

The returns of the Registrar General are not available for deducing conclusions regarding the incidence of Cancer owing to the want of accurate distinction between the different forms, and we are therefore practically confined to the use of hospital statistics which deal with cases selected as suitable for hospital treatment and are, therefore, of no value in determining the incidence of Cancer in the general population.

Bearing these difficulties in mind I propose to give a few results which can be obtained regarding the incidence of Cancer in respect to age, site, and sex, making use of the statistics compiled by the Imperial Cancer Research Fund from the General Hospitals of London (2 No. 2). Besides the error attached to hospital statistics mentioned above it must be remembered that these statistics are obtained from general hospitals and do not include cases from special hospitals such as hospitals for women and children. Hence the proportion of males to females does not represent the actual sex incidence and the statistics show an abnormally large number of cases of Carcinoma of the breast (1026) as compared with Carcinoma of the uterus (298). The statistics are, however, extremely valuable in that every case was verified microscopically by competent pathologists and hence valuable results can be obtained from a comparison of the figures *inter se*.

The statistics for Carcinoma deal with 1928 males and 2197 females and those for Sarcoma with 516 males and 391 females.

In comparing the carcinoma incidence in males and females we must remember that in females there are three situations which have no counterpart in males. The breast, uterus, and ovaries are, throughout a considerable period of life subject to periodical variations in the activity of cell proliferation and in



CHART 1.

the state of their blood supply. It is on this fact that the great liability of these organs to Carcinoma and other tumours depends. The breast, uterus, and ovaries together account for 1399 of the 2197 female cases of Carcinoma.

In the following charts which are compiled from these statistics the abscissæ represent the ages in quinquennial periods while the ordinates represent the number of cases per cent of the particular group of cases under consideration. For instance, the ordinates of the curve representing the incidence of Carcinoma in the male are calculated in percentages of the total male cases of Carcinoma and similarly for the other curves.

If we make a chart showing the age distribution of Carcinoma we obtain curves like those in Chart 1. One of these curves shows the distribution of the male cases, another of the female, and the third of the female cases omitting the breast, uterus, and ovaries.

All the curves are remarkably uniform and evidently follow a definite law. This will be found to be the case with all carcinoma curves which deal with a sufficient number of cases to eliminate fallacies.

We notice that all the curves commence to rise at about the age of 25, and, rising slowly at first and then rapidly, reach the maximum at 50 in females and 55 in males, subsequently falling rapidly. The effect of omitting the breast, uterus, and ovaries is to bring the female curve nearer to the position occupied by the male curve so that the maximum incidence occurs at the same age (55). The female curve is, however, still slightly in advance of the male curve.

We can conclude from these curves that age has an important influence on the incidence of Carcinoma.

If we plot out a similar curve for Sarcoma we find a marked difference (Chart 2). The curve starts at a moderately high point due largely to the inclusion of many blastocytomata. It descends rapidly to a minimum in the second and third age periods and then rises during the adolescent period. It subsequently remains at a high level with irregular fluctuations until it finally descends in old age.

It appears from these curves that age has not such a marked influence on the incidence of Sarcoma as it has in the case of Carcinoma. Omitting the initial high level due to congenital tumours, we see that the sarcoma curve rises rapidly during the adolescent period, while the carcinoma curve does not begin to rise markedly until after adult age has been reached.

If we study the age distribution of Sarcoma in different organs or groups of organs we notice several differences in the incidence. The curves, however, are not very reliable owing to the small number of cases in many situations. We find that if



CHART 2.

ve group together the glands—kidney, liver, pancreas, adrenal, nd parotid—32 per cent. of the cases occur in the first age eriod owing chiefly to the inclusion of the blastocytomata.

If, on the other hand, we take the breast, uterus, and ovaries we obtain a curve very similar to the carcinoma curve of the same organs but less regular.

Again, if we take the sarcomata of bone, we find that the curve rises rapidly during the adolescent period to a maximum



CHART 3.

at the age of 20—25, and then falls immediately to a level which is maintained almost constant until it falls at the age of 65. (Chart 3.)

Thus the age distribution in Sarcoma seems to follow no regular law. The one constant feature seems to be that the second and third age periods show the minimum incidence and that the incidence rises with adolescence.



CHART 4.

On the other hand, the curves of the age distribution of Carcinoma in different situations all show a remarkable resemblance to one another, rising rapidly, after adult age has been reached, to a maximum and then falling equally rapidly, the actual position of the maximum varying slightly in different cases.

If we compare the distribution of Carcinoma in the alimentary canal in males and females, we find that, as we should expect, the distribution is very similar in the two sexes, both curves showing identical variations. (Chart 4.) About the same proportion of cases in both sexes occurs in the stomach, but we notice that the male curve is higher in the parts above the stomach, while the female curve is the higher in the lower parts. While over 50 per cent. of the male cases occur above the stomach, over 50 per cent. of the female cases occur below.

This difference is presumably due to differences in the habits of the two sexes.

PART IV.

THE BEARINGS OF PATHOLOGY ON TREATMENT

ART IV. THE BEARINGS OF PATHOLOGY ON TREATMENT.

RECAPITULATION.

I propose now to consider the principles on which the treat-nt of Cancer should be based. First it will be necessary to apitulate the facts which we know about the disease.

1) Proliferation is an inherent property of the cell. Nor-lly this power of proliferation is kept in check by the influ-es of the organism as a whole exerted through adjacent cells.

(2) The essential causal factor in tumour formation is an rinsic one, namely the continued removal or diminution of the luences which restrain proliferation.

(3) This intrinsic factor may be of intrinsic origin alone or y arise from the action of extrinsic causal agents, as, for mple, in chronic irritation.

(4) The extrinsic causal factors are not of a specific kind are the agents we know generally as irritants.

(5) The extrinsic causal factors act partly by so influencing tissues as to give rise to the intrinsic conditions which I ve called *instability of equilibrium*. They also act as deter-ning factors in determining the appearance of a tumour when necessary intrinsic conditions are present.

(6) In either case the tumour itself at its first appearance is urely local condition.

(7) Certain chronic inflammatory conditions are sometimes lowed by cancer. Unfortunately we cannot say in any given se whether a given inflammatory condition will be followed Cancer or not. We cannot diagnose with certainty a pre-ncerous condition.

(8) Age has a marked influence in determining the incidence of Carcinoma but its influence in the case of Sarcoma is less marked.

(9) Tumours arise from normal tissues, from aberrant rudiments, or from pre-existing tumours.

(10) Tumours grow by proliferation of their own cells but the area of origin may show some extension. To this extent only do the surrounding tissues contribute to the growth of the tumour.

(11) Cancers extend mainly by infiltration of the surrounding parts, the proliferating cells penetrating the tissue spaces.

(12) The next stage of the process of extension is the permeation of lymphatics, veins, and other natural channels. In this process the cells grow continuously along the channels and the direction of the extension is not determined by the direction of the lymph or blood stream. During permeation of lymphatics in external cancers the extension is greater in the deep fascia than in the superjacent skin or subjacent muscle.

(13) By this process of permeation arise cutaneous nodules, invasion of bones, infection of the regional lymph glands, and, perhaps, invasion of the serous cavities.

(14) In the last stage of extension the cancer cells are detached and carried along by the lymph or blood stream or along mucous canals and give rise to metastatic deposits, *i.e.*, to deposits at a distance from the primary growth.

(15) Cancers, in some animals, can be transplanted to animals of the same species and cells displaced from a tumour can continue to proliferate.

(16) Tumours sometimes disappear spontaneously.

(17) In experimental cancer it is found that when a tumour has disappeared the animal subsequently shows some immunity. This immunity is specific, being most marked for the identical tumour from which the animal has recovered, less marked for other tumours of the same genus, and absent for tumours of another genus.

(18) Since the cells of which the tumours are composed are identical with normal cells it is probable that the metabolism in them is identical with the metabolism of the cells of origin.

aking allowance for the rapidity of proliferation in tumour lls.

(19) Cancers and other tumours are liable to the occurrence the same pathological processes as normal tissues. They n show inflammation, degenerations, necrosis, etc.

(20) Tumours in their growth give rise to an inflammatory action of the surrounding parts, resembling the reaction ainst foreign bodies, etc.

(21) While a simple tumour is to be regarded as an indivi- ual—a multicellular organism—a cancer is rather a colony of nicellular individuals.

(22) Cancer is truly described as an infection by cancer lls.

Principles of Treatment.

In estimating the value of any mode of treatment in the case Cancer two points must be borne in mind. The treatment ust be tried in a sufficiently large number of cases to eliminate llacies due to coincidence, etc., and all cases must be verified istologically by a competent pathologist.

We must also remember that cancers are often surrounded y an inflammatory zone which causes the dimensions of the umour to appear greater than the actual size. Hence an pparent improvement after treatment of any kind, as shown by diminution in size, may be entirely due to a diminution in the nflammatory zone with absorption of the products of inflamma- on while the actual tumour may not have been influenced at all.

It is much to be regretted that the writers and compilers of xt books of medicine seem to habitually ignore the subject of ancer. Cancer, however, has its medical aspects as well as its urgical, and we require more clinical observations on Cancer rom the medical point of view.

Since a tumour is a self-contained structure having in itself he power of growth, the most obvious method of treatment is o remove it entirely from the body. In this, the surgical treatment of Cancer, there are several points to be noticed. In the irst place cancer cells are themselves infective and if the wound e contaminated with them recurrence is likely to occur. In removing a cancer, then, care must be taken that the growth is not cut into during the operation. In the next place the growth infiltrates the surrounding tissues to a distance beyond that which can be estimated by microscopic examination. Hence the tumour must be removed with a sufficient margin of healthy tissue. Again Cancer spreads by permeation along lymphatic trunks and the extent of this permeation, in the case of external cancers, is greater in the deep fascia than in the superjacent or subjacent tissues. Hence the deep fascia should be removed to a greater extent than the skin. Mr. Handley, who has made a study of permeation in external cancers recommends that the deep fascia should be removed to an extent of two inches beyond the skin incision.[4] This permeation may extend to the regional lymph glands so that these should always be removed in continuity with the deep fascia and the primary tumour. The permeation also extends to an approximately equal extent in all directions from the primary growth and is not confined to the direction of the lymph current, so that the tissues should be removed to an equal distance all round the original tumour.

When metastasis has occurred the case may be regarded as beyond the limits of surgical operation.

Provided that the operation has included the removal of the whole tumour with its ramifications into the surrounding tissues this mode of treatment offers a prospect of complete cure.

Apart from removal, several attempts have been made to destroy the cancer cells *in situ* by caustics, etc. Unfortunately we do not know of any substance which has an injurious effect on the cancer cells which does not at the same time have a similar effect on the surrounding tissues. Among the caustics may be mentioned Potassium bichromate, and Potassium permanganate, both of which have been recommended. Another agent which has been recommended as having a special effect on cancer cells is Trypsin. This agent, about which we have heard so much, is supposed to act partly by its digestive properties and partly by the action of the associated amylopsin on glycogen.

Apart from attempting to destroy the cancer cells *in situ* we might attempt to starve them by cutting off their blood supply.

is, however, impossible to do this without at the same time arving the surrounding tissues, because the blood supply of a ncer is derived on all sides from the neighbouring parts. his method, then, is only applicable as a palliative method in re cases.

If we could detect a substance which was necessary for the etabolism of cancer cells to a greater extent than for the ormal tissues we might approach the treatment through this ubstance. Some people have thought that glycogen is such a ubstance, but recent researches have shown that glycogen is ot present in all tumours and that its presence does not depend n the malignancy so much as on the tissues from which the umour springs. It is obvious that if glycogen had been essenal for the proliferation of the cancer cells we might have tempted the treatment by taking steps for the elimination of e excess of the glycogen.

It is possible that some substance will yet be found through hich to approach the subject of treatment.

Mr. Holden Webb, on the basis of his hypothesis that ancer is due to the crystallisation of cholesterin out of the ving cells, suggested the use of hypodermic injections of soap olution with the idea of supplying a solvent for the cholesterin.[9]

Another possible manner of approaching the treatment of ancer would be to induce the tumour so to alter its mode of rowth as to behave as a simple tumour. The tumour would hen limit itself and so would be more amenable to surgical reatment. This idea is purely hypothetical as we have no nowledge of the influences which determine malignancy or on-malignancy.

Apart from any mode of treatment directed to the tumour tself we can conceive of modes of treatment directed towards ncreasing the resisting powers of the surrounding tissues. If ve could restore the influences normally existing between the omponent parts of the tissues involved, we should succeed in ausing the proliferation to cease and retrogression to set in. Here again we are met with the difficulty that we are at present otally ignorant of the nature of these influences, but some light progress has been made chiefly in connection with X rays and Radium, both of which appear to act by setting up an inflammatory reaction in the surrounding tissues combined with some destructive action on the tumour cells themselves. This method is most applicable in the case of cutaneous cancers, and the X ray treatment of Rodent ulcer and superficial squamous-celled carcinomata is admitted to be of considerable value.

Here also may be included suggested remedies such as Sodium cinnamate, and cumarate, which produce a general leucocytosis, also Coley's fluid which probably acts in a similar way.

Tonic treatment, especially arsenic, may be expected to raise the general resistance of the body tissues, and cases of amelioration under arsenic have been described. It is interesting to notice in this connection that arsenic has also a considerable beneficial effect on other diseases which are of a progressive nature, such as Hodgkin's disease, Leucocythæmia, Graves' disease, Diabetes insipidus, etc.

Another method which suggests itself is the general hygienic treatment including fresh air, exercise, freedom from worry, suitable diet, etc., which plays such an important part in improving the well-being of the body generally. We know that this method of treatment is proving of great value in infection by the tubercle bacillus and it is, perhaps, worth while to consider whether it would not have a similar effect in infection by cancer cells.

Another possibility in methods of treatment is in the use of extracts of glands which have an internal secretion. It is known that the pituitary, thyroid, and adrenals, have some effect on growth but what that effect is is not yet known. Thyroid treatment has been tried in some cases and at the Christie Hospital we have tried pituitary extract in two cases but without result. Thymus extract has also been tried. Removal of the ovaries and testicles has been tried in some cases of inoperable cancer with apparent benefit in some cases, but we must not forget that in the opinion of some authorities the removal of the ovaries increases the liability to Cancer.

Lastly we have empirical methods of treatment which may be described as modes of treatment tried hap-hazard without any scientific basis. Such methods are violet leaves, molasses, chian turpentine, etc. We may say that no substance has yet been found that has any effect on Cancer, nor is it likely that any substance will ever be found which has any effect comparable with that of quinine in Malaria since Malaria is due to an extrinsic agent while Cancer is, as we have seen, an infection by cells derived from the body cells, and it is hardly possible to conceive a substance which will destroy the tumour cells without at the same time destroying their near relations, the body cells.

With regard to the effects of these different modes of treatment we can say that, apart from surgical operation, and, in special cases, X-rays or radium, none of the numerous suggested remedies have proved of any permanent value. Cases of amelioration during treatment of various kinds have been described, but further investigation has always shown that these cases of improvement are accidental and cannot be ascribed to the immediate effect of the treatment. Some of these cases of improvement are due to the lessening of the pericancerous inflammation producing an apparent diminution in the size of the tumour.

When the case is operated upon early enough to allow of complete removal we have a fair prospect of complete cure. When, however, it has gone beyond the limits of surgical operation the prospects are at present almost hopeless. Surgery and medicine can still do much in the way of palliative measures but we know of no remedy which will effect a cure.

I may say that all suggested methods of treatment, whatever their origin, are given a fair trial at the Christie Hospital provided that the use of them does not involve danger to the patient.

It appears most reasonable, in considering possible modes of treatment, to direct our attention more towards improving the general condition of the patient so that his tissues may resist the growth of the tumour, rather than to attempting to find a substance which will destroy the cancer cells without destroying the other tissues. It is possible, also, that if we can discover a chemical substance which is necessary for the proliferation of the cancer cells, we may be able to approach the treatment of Cancer through this substance.

Whether or no we shall ever see a treatment by immunisation remains to be seen. I have mentioned that mice which have recovered from Cancer are immune to that particular cancer and, to a less degree, to other related forms, and, if this mode of treatment ever becomes possible, it would seem probable that it will consist in immunising a patient against his own cancer.

With regard to preventive treatment the only means that can be suggested are ordinary hygienic principles and the prompt treatment of any chronic inflammatory foci that may occur.

CONCLUSION.

I have now said what I have to say about Cancer. I have considered the pathology of the disease and have indicated the bearing of the facts of pathology on the indications for treatment. I conclude by reminding you that the problem of Cancer is the problem of Growth. When biologists can tell us what are the influences which determine the form and size of the living organism and of its constituent parts, then pathologists will be in a position to determine more exactly the nature of tumours. In the meantime all features in tumours must be investigated and such investigation must not be limited to tumours alone. Other diseases and normal processes must be included in the scope of the investigation, as also problems of chemistry and physics, before we can arrive at a complete answer to the question "What is Cancer?"

APPENDIX

APPENDIX.

THE PARASITIC THEORY.

I have not thought it necessary in the lectures to enter into
full discussion of this theory as I went into the matter fully
my Erasmus Wilson Lectures in 1902.[10] The criticisms I
en brought forward have never been answered, and it is a
rious fact that the upholders of this theory never think it
cessary to answer the criticisms brought against it, while they
ntinue to bring forward again and again the same arguments
its favour. Very few of the leading pathologists have ex-
essed an opinion in favour of the theory that the essential
sual agent of Cancer is a parasite. The upholders of the
eory are mostly surgeons or bacteriologists who do not appre-
ate the pathological and biological difficulties which the theory
volves.

For the sake of those who are not well acquainted with the
scussion which has ranged about this theory I here introduce
brief résumé of the argument.

The grounds on which the theory is based are :—

(1) *The apparent resemblance of Cancer to such infective*
seases as Tuberculosis, etc. I have considered this point in
e lectures and I have pointed out that the only argument to
obtained from this resemblance is the argument that the
ncer cells themselves are parasites.

(2) *The inoculability of Cancer.* If Cancer were due to a
rasite we should expect infection of one person by another to
of frequent occurrence. Such cases are, however, extremely
re, and hardly more than could be explained by coincidence.
he venereal growths which are met with in dogs are highly
ntagious, but it is not certain whether they are to be considered
true tumours or whether they are more allied to granulomata.
icroscopically they show a close resemblance to sarcomata but
ey do not show the same power of infiltrating the surrounding
ssues. Even if they be considered as sarcomata the fact that
ey are contagious is no argument in favour of their being of

parasitic origin, since the infecting agents may be the tumour cells themselves.

In the transmission of Cancer from mouse to mouse the new tumour arising is a direct continuation of the growth of the introduced portion and is not due to infection of the tissues of the new host.

(3) *The lesions produced in the rabbit's liver by the coccidium oviforme.* In coccidiosis we find local dilatations of the bile ducts filled with coccidia and epithelial debris. The younger forms of the parasite are contained within the epithelial cells while the older lie free in the cavity. In no case do we find any lesions at all resembling Cancer. The process appears to be of the nature of a catarrhal inflammation, the infected epithelial cells being cast off and the remaining cells proliferating to take their place. The coccidia are sometimes found in the connective tissue where they give rise to an ordinary chronic inflammation of the fibrous tissue.

(4) *The topographical distribution of Cancer.* Cancer has been supposed to occur endemically in certain localities or certain houses. The evidence, however, is not strong on this point nor are there any resemblances between the circumstances of different cancer districts. Some cancer districts are in valleys, others on hills; some are in moist situations, others in dry deserts, etc. It has been proved that the cells of a mouse cancer can retain their vitality for a long time outside the body under various circumstances and yet be able to proliferate on being introduced into a new mouse. Hence the endemic occurrence of Cancer is in itself no argument for the necessity of a parasitic causal agent.

(5) *The presence of certain cell inclusions, supposed to be of a parasitic nature and the possibility of obtaining cultures of bacteria, etc., from cancers.* It does not appear that any of the numerous observers in this field agree with others as to what the cancer parasite is. Bacilli, cocci, protozoa, blastomycetes, and moulds have all had their day. Each new observer points out the mistakes made by his predecessors and then brings forward a new parasite to be, in its turn, demolished by his successors. The inclusions described as parasites include various forms of

ıerative changes (hyaline, colloid, vacuolation, etc.), archo-
ıic vesicles and centrosomes, extruded nucleoli, detached
ɔns of nuclear chromatin, and, in all probability, globules
ıolesterin. As regards cultivation experiments the same
ılties arise. There is no doubt that organisms can be culti-
from cancers in some cases, but there is no evidence that
have anything to do with the causation of Cancer. A large
ɔer of different bacteria and blastomycetes have been culti-
at different times, but no one has yet succeeded in repro-
.g undoubted Cancer on inoculating animals with the
res of the suspected organisms.

hus the grounds on which the theory is based are wholly
quate. There are numerous other objections to the
y.

) *It is entirely unnecessary.* All the phenomena of
er can be explained on the supposition that the cancer cells
hemselves the infecting agents. As I have pointed out,
apacity for proliferation is an inherent property of the cell
is of all living organisms and it requires no parasitic theory
plain it.

) *The theory will not explain the origin of Cancer.* We
of no parasites which are capable of producing a primary
feration of cells. The first effect of any irritant is destruc-
ɔr injury to the cells. There is no evidence that in Cancer
is any initial injury. Even if we admit the existence of a
ite which produces a primary cell proliferation this would
xplain Carcinoma, since the essential feature in Carcinoma
netration of the epithelial cells into the connective tissue
s.

) *The theory will not explain the growth of Cancer.* The
ion against any irritant, *e.g.*, Bacillus tuberculosis, is a
ipetal proliferation of the tissue cells towards the irritant,
eas in Cancer the growth is centrifugal into the surround-
issues.

) *It will not explain the site incidence of Cancer.* For
ple, it will not explain why the small intestine is so rarely
eat of primary Carcinoma, nor why the acini of the breast
nore frequently affected than the ducts, nor will it explain
the breast and uterus are more frequently affected than
organs. In the case of Sarcoma we should have to

assume that the parasites are conveyed to the site of origin by the blood, and in that case we should expect to find that multiple primary growths were more common than they are.

(10) *It will not explain why there is an almost complete immunity to Carcinoma in the first twenty years of life.*

(11) *It will not explain the congenital tumours.*

(12) *It will not explain the occurrence of mixed tumours, such as the chondrosarcomata, etc., and especially the blastocytomata.*

(13) If all forms of Cancer are supposed to be due to the same parasite it is impossible to explain why metastatic growths always resemble the primary. We should expect, for instance, that a carcinoma would sometimes give rise to sarcomatous metastatic growths, and we should also expect that, if a carcinoma of the breast gave rise to a metastatic nodule in the liver, the latter would show the characteristics of a hepatic carcinoma, etc.

If, on the other hand, each kind of cancer has its own parasite we are met with the difficulty of the infinite variety of cancers. We should have to assume different parasites for all the different species of Carcinoma and Sarcoma, and we should have to imagine that each organ and tissue in the body has its own series of cancer parasites, and, moreover, that each species of animal has a different series of cancer parasites to every other animal.

(14) It is not usually contended that simple tumours are due to parasites, but, if they are due to extrinsic casual agents these agents must be of a parasitic nature since a continued and increasing growth demands a continued and increasing causal agent, if any agent is considered necessary. If it be contended that simple tumours are not due to extrinsic agents, then I answer that there is no justification whatever for separating the malignant tumours from the simple, and to say that while the simple are due to intrinsic factors alone, the malignant are due to the action of parasites.

We see then that the parasitic theory will not explain any of the phenomena, and that it vastly increases the difficulty of explanation.

We can say, then, that Cancer is not due to a specific parasite or parasites, but, on the other hand, we can say that the cancer cells themselves act as parasites. This latter view will explain all the phenomena of Cancer.

Fig. 1.



Tissue Tumour. Adenoma of the Liver.

This is composed of cells, resembling hepatic cells, arranged in columns as in the normal liver. The tumour from which the section was taken was about 8 inches in diameter and was surrounded by a capsule. Throughout the whole tumour there was no arrangement of the cell columns into lobules and no ducts were found in any part of the growth.

Fig. 3.



Tissue Tumour. Chondroma.

The nodules of cartilage, of which this tumour is composed, are everywhere surrounded by fibrous tissue.

Fig. 4.



Cell Tumour. Chondrosarcoma.

The nodules of cartilage are not sharply marked off from the surrounding sarcomatous tissue, but there is a gradual transition from one to the other, the sarcoma cells themselves becoming chondrocytes, thus giving rise to the cartilage as a secondary product.

Fig. 5.



Tissue Tumour. Adenoma of the Rectum.

The epicytes are everywhere arranged in definite tubes and sharply marked off from the surrounding stroma. There is no infiltration.

Fig. 6.



Cell Tumour. Carcinoma of the Rectum.

In this part of the tumour the structure somewhat resembles that of an adenoma. The tubes are, however, more irregular and in some places they are lined by several layers of cells.

Fig. 7.



Cell Tumour. Carcinoma of the Rectum. From the same section as Fig. 6.

This shows solid columns of epicytes penetrating the surrounding connective tissue, the tubes being of secondary formation.

Fig. 8.



Cell Tumour. Carcinoma of Rectum. From the same section as Figs. 6 and 7.

This is a portion of Fig. 7 photographed with a higher power. It shows the formation of a lumen in a solid column of epicytes. Above are seen two other solid columns of cells penetrating the connective tissue.

Fig. 9.



Angeioma of the Liver. (From Dr. Mantle's specimen.)

There are groups of hepatic cells in the septa between the cavernous spaces.

Fig. 10.



Blastocytoma of the Kidney. Low power.

The section shows groups of small round cells in which are embedded epithelial tubes. There are also strands of spindle cells running between the groups of round cells. Other parts of the tumour contained mucous tissue and smooth and striated myocytes.

Fig. 11.



Blastocytoma of the Kidney. High power.

This shows the earliest stage of the formation of epithelial tubes from the round cells. In the upper part of the section is seen a group of the small round fundamental cells. Below this is another group slightly more advanced showing the commencement of the formation of a lumen.

Fig. 12.



Blastocytoma of the Kidney. High power.

This shows a later stage in the formation of epithelial tubes. In the centre of the section is seen a group of cells with epithelial characters surrounding a lumen which contains débris and a detached cell.

Fig. 13.



Blastocytoma of the Kidney. High power.

This shows a space lined with epicytes surrounded by mucous tissue. To the left is a group of the fundamental cells. There is no sharp demarkation between the epithelial cells, the mucous tissue cells, and the fundamental cells, all being connected together by cell-processes.

Fig. 14.



Blastocytoma of the Kidney. High power.

This section shows the invagination of one side of an epithelial tube to form a glomerulus-like structure,

Fig. 15.



Blastocytoma. Parotid Tumour.

This section shows several groups of cells, embedded in which are epithelial tubes, some of which contain gelatinous material. The groups of cells are separated by areas of mucous tissue. There is no sharp demarkation between the different kinds of cells, the epithelial cells and the mucous tissue cells both passing gradually into the intermediate fundamental cells.

Fig. 16.



Primary Carcinoma of the Femur.

The cells in this tumour are of the epithelial type and, in places, are typically columnar in shape. There were no growths elsewhere in the body.

Fig. 17.



Primary Carcinoma of the Pleura. From a child aged 2.

This is a typical spheroidal-celled carcinoma.

Fig. 18.



Primary Carcinoma of a Rib.

This shows typical squamous cells with prickles.

Fig. 19.



Sarcoma of the Neck. From a guinea pig.

This formed a tumour the size of a marble in the subcutaneous tissue of the neck. There was a similar growth in the neighbourhood of the pancreas.

Fig. 20.



Carcinoma of the thyroid. From a mouse.

The dilated vesicles of the thyroid body are seen on the right of the photograph, the carcinoma being to the left. There were secondary tumours in both axillæ. Some of the tubes in the primary and secondary tumours contained some colloid secretion.

Fig. 21.



Lipoma. From a parrot.

This was a large tumour in the subcutaneous tissue covering the pectoral muscles. It shows the typical structure of a lipoma.

Fig. 22.



Giant-celled Sarcoma of the Femur. From a drake.

This is a typical giant-celled sarcoma. There were secondary growths in the liver.

Fig. 23.



Carcinoma of the Kidney. From a rabbit.

There are several nodules of growth in the upper pole of the kidney.

Fig. 24.



Carcinoma of the Kidney. From a rabbit.

This is a photograph of a section of the tumour shown in Fig. 23. It is a columnar-celled carcinoma.

Fig. 25.



Carcinoma of the Breast, infiltrating the Pectoral Muscle.

This shows the carcinoma cells penetrating between the muscle fibres, causing them to undergo atrophy *in situ*.

Fig. 26.



Carcinoma of the Breast invading fat.

The carcinoma cells are proliferating within the fat cells.

Fig. 27.



Carcinoma infiltrating the wall of a large vein and forming growth in the intima.

Fig. 28.



Carcinoma of the Breast. Polarised light.

The section has been heated and allowed to cool. Numerous anisotropic globules are seen appearing as bright globules divided into four parts by a dark cross. These were lying in and among the carcinoma cells which are not seen by this method of illumination.

Fig. 29.



Normal Adrenal body. Polarised light. Low power.

The cells of the cortex are filled with numerous crystals appearing bright on a dark background.

Fig. 30.



Normal Adrenal body. Polarised light. High power.

This is from the same section as Fig. 29 and shows the characters of the crystals in the adrenal cortex.

Fig. 31.



Normal Adrenal body. Polarised light. High power.

This is from the same section as Figs. 29, 30. The section has been heated and allowed to cool and the photograph shows the presence of anisotropic globules instead of crystals. Only one globule is in focus. Compare with Fig. 28.

Fig. 32.



Bilharzial Carcinoma of the Bladder. (From Dr. Harris' specimen.)

This section shows a typical carcinoma of the bladder. No ova were found in the tumour itself.

Fig. 33.



Bilharzial Carcinoma of the Bladder. (From Dr. Harris' specimen.)

This section is from the connective tissue in the neighbourhood of the tumour and shows numerous ova.

REFERENCES.

Bashford (E. F.). "Liquid Air and Cancer." *Lancet*, 1908, vol. i, p. 394.

Bashford (E. F.) and Murray (J. A.). "Scientific Reports of the Imperial Cancer Research Fund."

Beard (J.). "Embryological Aspects and Etiology of Carcinoma." *Lancet*, 1902, vol. i, p. 1706.

Handley (W. Sampson). "Third, Fourth and Fifth Reports of the Cancer Research Laboratories of the Middlesex Hospital."

Harris (T.). "Bilharzia Hæmatobia co-existing with an Epitheliomatous Growth in the Bladder." *Trans. Path. Soc.*, vol. xxxix, 1888.

Mantle (A.). "An unusually large Angeioma of the Liver." *Brit. Med. Journ.*, 1903, vol. i, p. 365.

Ribbert (H.). "Geschwulstlehre." Bonn, 1904.

Salvin-Moore (J. S.) and Barratt (J. O. Wakelin). "Note on the effect of Liquid Air on the Graftable Cancer of Mice." *Lancet*, 1908, vol. i, p. 227.

Webb (J. Holden). "Cancer, its Nature and Treatment." *Lancet*, 1901, vol. ii, p. 976.

White (C. Powell). "The General Pathology of Tumours." *Lancet*, 1902, vol. i.

White (C. Powell). "Cholesterol, Fluid Crystals and Myelin Forms." *Medical Chronicle*, vol. xiv, fourth series, 1908.

White (C. Powell). "Crystals in Tumours." *Journal of Path. and Bact.*, vol. xiii.

White (C. Powell). "The so-called Fatty Degeneration of the Adrenals." *Ibid.*

White (C. Powell). "On the Relation between Cause and Effect in Disease." *Brit. Med. Journal*, 1900, vol. ii.

2980

INDEX

INDEX.

PAGE
Aberrant rudiments 12
Adenocarcinoma 20
Adenoma 5, 10, 46
Adenomata 8
Adrenals, crystals in 44
Adrenals, relation of, to growth and development 45
Age distribution of carcinoma 52
—— —— sarcoma 52
Age, influence of 49
Alimentary canal, distribution of carcinoma in 56
Anæmia, pernicious 9
Angeioma of liver, growth of 14
Animals, cancer in 20
Anisotropic globules 43
Area of origin, extension of 14, 21
Arsenic treatment 64
Arthritis, rheumatoid 9
Atrophy, progressive 8

Beard's hypothesis 12
Bilharzial carcinoma 35, 48
Blastocytes 17
Blastocytoma of kidney 17
—— parotid 18
Blastocytomata 17
Blood-vessels in cancer 27
Bones, infiltration of 23

Cachexia 30
Cancer, analogy between tuberculosis and 34
—— in animals 20
—— area of origin 21
—— definiton 3
—— destructive characters of 22
—— apparent encapsulation of 22
—— experimental 21
—— gall-stones in 44
—— growth of 22, 40, 42
—— immunity in 28
—— infiltration of tissues 22
—— influence of civilisation on 49

PAGE
Cancer, inoculability ... 69
—— in mice ... 21, 27, 28, 41, 70
—— micro-organisms in ... 70
—— penetration of fat cells ... 22
—— penetration of muscle fibres ... 22
—— rudiment of origin ... 21
—— topographical distribution of ... 70
—— cells, effect of cold on ... 41
—— —— conjugation in ... 41
—— —— resemblance to protozoa ... 41
Carcinoma ... 5, 46
—— age distribution of ... 52
—— bilharzial ... 48
—— distribution of, in the alimentary canal ... 56
—— growth of ... 10
—— sarcoma arising from ... 25, 27
—— stroma in ... 26
Carcinomata ... 17, 19
Carcinosis, diffuse ... 13, 21
Causal factors ... 33
—— —— adjuvant ... 33
—— —— essential ... 33
—— —— extrinsic ... 33, 34
—— —— intrinsic ... 33, 35
Caustics ... 62
Cell inclusions ... 70
—— proliferation ... 35
—— tumours ... 5
Cells ... 4
—— indifferent ... 17
—— influence of, on stroma ... 26
—— nomenclature of ... 7
Central growth ... 14
Centrifugal growth ... 14
Centripetal growth ... 14
Cholesterin in adrenals ... 44
—— in tumours ... 43
Chondroma, growth of ... 10
—— metastasis in ... 24
Chondro-sarcoma, growth of ... 10
Ciliary movement, co-ordination of ... 38
Cirrhosis of the liver ... 8
Civilisation, influence of, on cancer ... 49
Classification ... 5, 6
—— basis of ... 7
—— embryological ... 7
Clinical features of cancer ... 29
Coccidiosis ... 70
Cohnheim's theory ... 11

PAGE
Cold, effect of, on cancer cells 41
Coley's fluid 64
Conjugation in cancer cells 41
Co-ordinating mechanism of the organism 37
Co-ordination of ciliary movement 38
Crystals in adrenals 44
—— in tumours 43
Cutaneous nodules 29
Cytomata 5, 17
—— distinction between histiomata and 5, 10, 11

Definition 3
Degenerative processes 28
Destructive characters of cancer 22
Dermoid cyst 4
Destination of cancers 27
Development 39
Diabetes insipidus 9
—— mellitus 9
Diagnosis, histological, of malignancy 11
Difference between simple and malignant tumours 40
Differentiation 37
—— in blastocytomata 19
Diffuse carcinosis 13, 21
Disappearance, spontaneous, of cancers 28
Dogs, venereal growths in 69
Encapsulation, apparent, of cancers 22
Endothelial tumours 19
Endothelioma 8
Epicytes 17
Epithelial cells 17
—— pearl 46
Epithelioma 8
Equilibrium, physiological 37, 45
—— stable 46
—— unstable 46, 47
Erythrocythæmia 9
Exophthalmic goitre 9
Expansive growth 11, 14
Experimental study of cancer 21
Extension, modes of 23
Extrinsic causal factors 33, 34

Fat cells, penetration of 22
Fibroma 4
Fibrosis, progressive 8

INDEX

PAGE
Gall-stones in cancer 44
Glycogen in tumours 29, 63
Goitre 8
—— exophthalmic 9
Growth 39
—— centrifugal 14
—— centripetal 14
—— expansive 11, 14
—— infiltrative 11, 14
—— peripheral 14, 22
—— adenoma 10
—— angeioma 14
—— cancer 22, 40, 42
—— carcinoma 10
—— chondroma 10
—— chondra-sarcoma 10
—— cytomata and histiomata 11
—— simple tumour 39
—— of tumours 13
Growths, venereal, of dogs 69

Habit of growth 47
Heredity 48
Histiomata 5
—— nomenclature of 8
—— distinction between cytomata and 5, 10, 11
Hodgkin's disease 9
Hygienic treatment 64
Hyperergasis 9, 47
Hypertrophy, progressive 8
—— of the prostate 8
Hypo-ergasis 9
Hypothesis, Beard's 12

Immunity in cancer 28, 66
Indifferent cells 17
Infiltration a sign of malignancy 10, 11
—— of tissues 22, 23
Infiltrative growth 11, 14, 22
Inflammatory conditions, influence of 48
Influences, intercellular 38, 42
—— limiting cell proliferation 37
Inoculability of cancer........ 69
Intercellular bridges 38, 39
—— influences 38, 42
Intrinsic causal factors 33, 35

Kidney, mixed tumour 17

PAGE
Leucocythæmia 9
Leucoplakia 48
Liver, cirrhosis 8
Living organisms, characters of 35
Lupus cancer 35, 48

Malignancy, histological diagnosis of 11
—— infiltration a sign of 11
—— signs of 10
Malignant tumours 7
—— difference between simple and 9
Mesocytes 17
Metabolism in cancer 29
Metaplasia 12
—— progressive 9
Metastasis 23
—— in apparently simple tumours 24
—— in chondromata 24
Mice, cancer in 21, 26, 27, 28, 41, 70
Micro-organisms in cancer 70
Mitosis 42
Mixed tumour of the kidney 17
—— —— parotid 18
Multicellular organism, nature of 38
Multiple tumours 25
Muscle fibres, penetration of 22
Muscles, infiltration of 23
Muscular atrophy, progressive 8
Myositis ossificans 9

Nomenclature 7

Organism, co-ordinating mechanism of 37
—— relation of tumour to 39
—— multicellular, nature of 38
Organomata 5
Organs 4
Organ tumours 5
Origin, area of 14, 21
—— rudiment of 11, 13, 21
—— sites of 36
Ovariotomy for cancer 64

Papillo-carcinoma 20
Papilloma 8, 46
Parasitic theory 34, 69
Parotid tumour 18
Penetration of fat cells and muscle fibres 22
Peripheral growth 14, 22

PAGE
Permeation 23
Pernicious anæmia 9
Physiological equilibrium 45
Pituitary treatment 64
Processes, progressive 9
—— reactive 9
—— resembling tumour growth 8
Progressive hypertrophy 8
—— processes 9
Proliferation 35, 36
—— influences which limit 37
Prostate, hypertrophy of 8
Protozoa, resemblance between cancer cells and 41

Radium 64
Reactive processes 9
Recapitulation 59
Recurrence 30
Regeneration and repair 46
Relation of the tumour to the organism 39
Rheumatoid arthritis 9
Ribbert's theory 13
Rodent ulcer 22
Rudiment, Cohnheim's 11, 13, 19
—— of origin 11, 13, 19, 21
Rudiments, aberrant 12

Sarcoma 4, 47
—— age distribution of 52
—— arising from carcinoma 25, 27
Sarcomata 17, 19
Sex, influence of 49
Signs of malignancy 10
Simple tumours 7
—— —— growth of 39
—— —— metastasis in 24
Simple and malignant tumours, difference between 9
Sites of origin 36, 49
Skin, infiltration of 23
Soap treatment 63
Sodium cumarate 64
Spontaneous disappearance of tumours 28
Statistics 49
Stroma in carcinoma 26
—— influence of cells on........ 26
Surgical treatment 61

Theory, the parasitic 34, 69
Thyroid treatment 64

PAGE
Tissue tumours 5
Tissues 4
—— of secondary formation 10, 11, 19
Topographical distribtuion of cancer 70
Treatment, arsenic 64
—— caustics 62
—— Coley's fluid 64
—— ductless glands 64
—— empirical 64
—— hygienic 64
—— immunisation 66
—— ovariotomy 64
—— preventive 66
—— principles of 61
—— radium 64
—— soap 63
—— sodium cumarate 64
—— surgical 61
—— trypsin 62
—— X-rays 63
Trypsin 62
Tuberculosis, analogy between cancer and 34
Tumour, definition 3
—— mixed, of parotid 18
—— relation of, to the organism 39
—— rudiment of origin 11
Tumour growth, relation of, to other processes 8
Tumours, cell 5
—— crystals in 43
—— difference between simple and malignant 9, 40
—— endothelial 19
—— glycogen in 29
—— growth of 5
—— metastatic 24
—— multiple 25
—— organ 5
—— sites of origin 36
—— tissue 5

Ulcer, rodent 22

Venereal growths in dogs 69
Vital energy 35, 43

X-rays 63

Publications of the University
of Manchester

MANCHESTER UNIVERSITY PUBLICATIONS.

ANATOMICAL SERIES.

No. I. STUDIES IN ANATOMY from the Anatomical Department of the University of Manchester. Vol. iii. Edited by ALFRED H. YOUNG, M.B. (Edin.), F.R.C.S., Professor of Anatomy. Demy 8vo, pp. ix, 289, 23 plates. 10s. net. (Publication No. 10, 1906.)

"All the papers contained in the volume are real additions to the knowledge of the subject with which they deal. For three of the studies Prof. Young is either in part or wholly responsible, and he is to be congratulated on the vigour shown by the Manchester School of Anatomists."—*Nature.*

"This work affords admirable evidence of the virility of our younger British Universities. It is a notable addition to an already notable series."—*Medical Review.*

"This forms the third volume of the Studies in Anatomy issued by the Council, and contains contributions of considerable interest. The volume is well printed and bound. It speaks well for the activity of investigation at Manchester."—*Lancet.*

"The volume is well got up and is evidence of the continuation of the excellent work which has been carried on for so long a period, under Professor A. H. Young's supervision, and has been encouraged and stimulated by his own work."—*British Medical Journal.*

"Throughout the papers, careful research and accurate observation are manifested, and they will repay careful perusal. To the Anatomist, as well as the practical physician or surgeon, they will prove valuable."
—*Edinburgh Medical Journal.*

CLASSICAL SERIES.

No. I. A STUDY OF THE BACCHAE OF EURIPIDES. By G. NORWOOD, M.A., Assistant Lecturer in Classics. Demy 8vo, pp. xx, 188. 5s. net. (Publication No. 31, 1908.)

"The interest of Mr. Norwood's book, which . . . is a very welcome addition to the bibliography of Euripides, and a scholarly and interesting piece of work, displaying erudition and insight beyond the ordinary, lies in the way in which, by applying Dr. Verrall's methods . . . . he first shows up difficulties and inconsistencies, some of which have hardly been noticed before . . . . and then produces his own startling theory, which he claims is the great solvent of all the perplexities."
—*Saturday Review.*

"Unless very strong evidence can be produced against Mr. Norwood's view, it must be accepted as the true solution of the problem. . . . Mr. Norwood is generally clear, and abounds in illuminating thoughts. He has added a full bibliography (running to twenty-three pages) of writings on Euripides, and for this every scholar will offer his sincere thanks. . . . He has done a very good piece of work."—*Athenæum.*

"This volume forms the first of a Classical Series projected by the Manchester University, who are to be congratulated on having begun with a book so original and full of interest. . . . It is admirably argued, and is instinct with a sympathetic imagination. It is, at the very least, an extremely able attempt to solve a very complex problem."
—*Manchester Guardian.*

"Mr. Norwood demonstrates on every page his scholarship and knowledge, and gives proof of much painstaking research. The treatise is as valuable as it is interesting."—*Manchester City News.*

"It is a most ingenious theory, and a reviewer whom it has left unconvinced is all the more bound to give his testimony to the consistent skill, learning, and independence of judgment with which it is presented. The book . . . strikes us as the product of vigorous and independent thought."—*Times.*

MANCHESTER UNIVERSITY PUBLICATIONS.

ECONOMIC SERIES.

No. I. THE LANCASHIRE COTTON INDUSTRY. By S. J. CHAPMAN, M.A., M. Com., Stanley Jevons Professor of Political Economy and Dean of the Faculty of Commerce. Demy 8vo, pp. vii. 30. 7s. 6d. net. (Publication No. 4, 1904.)

"Such a book as this ought to be, and will be, read far beyond the bounds of the trade."—*Manchester Guardian.*

"There have been books dealing with various phases of the subject, but no other has so ably treated it from the economic as well as from the historical point of view."—*Manchester Courier.*

"The story of the evolution of the industry from small and insignificant beginnings up to its present imposing proportions and highly developed and specialised forms, is told in a way to rivet the attention of the reader . . . . . . the book is a valuable and instructive treatise on a fascinating yet important subject."—*Cotton Factory Times.*

"Highly valuable to all close students."—*Scotsman.*

(GARTSIDE REPORT, No. 1.)

No. II. COTTON SPINNING AND MANUFACTURING IN THE UNITED STATES OF AMERICA. By T. W. UTTLEY, B.A., Gartside Scholar. Demy 8vo, pp. xii. 70. 1s. net. (Publication No. 8, 1905.)

"Mr. Uttley is to be congratulated on the performance of a not altogether easy task, and his book, in conception and execution, appears to fulfil admirably the intentions of the Trust."—*Manchester Courier.*

"The writer gives ample details concerning wages and other features connected with typical mills . . . and the information thus gathered is of interest and value to the factory operative as well as the student and economist."—*Cotton Factory Times.*

"Mr. Uttley describes how he visited the mills in various States in a very systematic and detailed manner. Altogether the report makes an admirable and welcome collection of information, and will be found on many occasions worthy of reference."—*Textile Mercury.*

(GARTSIDE REPORT, No. 2.)

No. III. SOME MODERN CONDITIONS AND RECENT DEVELOPMENTS IN IRON AND STEEL PRODUCTION IN AMERICA, being a Report to the Gartside Electors, on the results of a Tour in the U.S.A. By FRANK POPPLEWELL, B.Sc., Gartside Scholar. Demy 8vo, pp. vi. 119. 1s. net. (Publication No. 21, 1906.)

"The American methods of iron and steel production are described, from the practical as well as the statistical side."—*Manchester Courier.*

"Mr. Popplewell writes clearly and well, and he is to be congratulated upon having carried his task through in so entirely a satisfactory manner."—*Manchester City News.*

"America's progress in iron and steel is more wonderful than any bald statistics of production with which we are so familiar can indicate. How that progress has been effected—effected under labour, transport and other difficulties—Mr. Popplewell tells us in an interesting and keenly intelligent review."—*Manchester Guardian.*

"A minute observation of detail . . . characterises the whole work."
—*Iron and Coal Trades Review.*

"Mr. Popplewell gives a clear exposition of the results of specialisation in production, of the development of ore-handling machinery, and of the general use of the charging machine, features that characterise American practice. He shows, too, that the colossal blast-furnace with huge yield due to high-blast pressure, regardless of consumption of steam and boiler coal, is giving place to a blast furnace of more modest dimensions. . . .

MANCHESTER UNIVERSITY PUBLICATIONS.

ECONOMIC SERIES.

"The impression derived from reading Mr. Popplewell's report is that many of the most striking developments, admirable as they are, were designed to meet special wants, and are not necessarily applicable in Great Britain."—*Nature.*

"The book has its interest for the educationist as well as for the manufacturer."—*Scotsman.*

"A chapter of special interest to British consumers is one devoted to the consideration of raw materials."—*Glasgow Herald.*

(Gartside Report, No. 3.)

No. IV. ENGINEERING AND INDUSTRIAL CONDITIONS IN THE UNITED STATES. By Frank Foster, M.Sc., Gartside Scholar. Demy 8vo, pp. ix. 106. 1s. net.

(Publication No. 22, 1906.)

"The report under review is of very great interest to those connected with the manufacturing branch of engineering in this country, many of whom will have to relinquish their preconceived notions regarding American methods, if Mr. Foster's conclusions are to be accepted."

—*Electrical Review.*

"The book altogether is very readable, and one we can heartily recommend to all interested in the economics of engineering."

—*The Practical Engineer.*

"Mr. Foster's observation of facts is fresh and interesting . . . . the technical side of his report exhibits much care."—*Manchester Guardian.*

"The book is well worth reading."—*Iron and Coal Trades Review.*

"There is much in the book which will be new to English readers, even to those who have studied the reports of the Moseley and other recent 'commissions.'"—*Belfast News Letter.*

No. V. THE RATING OF LAND VALUES. By J.D. Chorlton, M.Sc. Demy 8vo, pp. viii. 177. 3s. 6d. net. (Publication No. 23, 1907.)

"A timely and temperate treatise on a subject of growing interest.

—*Pall Mall Gazette.*

"The writer is learned, intelligent, progressive, fair and lucid."

—*Progress.*

"The facts and deductions are well put."—*Western Mail.*

"Chapters upon the scheme of the Royal Commission (minority report) —'Building Land,' 'The Future Increase of Land Values,' 'The Municipal Bill,' and others . . . set forth with clearness and detail some of the many interesting and difficult subjects in connection with valuation, rates and rating."—*Estates Gazette.*

"Mr. Chorlton has made a contribution to this interesting controversy which is worthy of the serious attention of all persons interested in the subject."—*Local Government Chronicle.*

"The arguments for and against this proposed reform in the taxation of land have never been more fairly and freely stated."

—*Liverpool Daily Post and Mercury.*

"Mr. Chorlton deals clearly and concisely with the whole subject of rating and land values."—*The Standard.*

"The impartiality and candour of Mr. Chorlton's method are beyond dispute, and his book will repay careful study by all who are interested in the question, from whatever motive."—*Westminster Gazette.*

"The first half of this book deserves to become a classic . . . . . . is one of the best books on a practical economic question that has appeared for many years. It is not only scientifically valuable, but so well written as to be interesting to a novice on the subject."—*The Nation.*

"This thoughtful and judicially expressed treatise."

—*Manchester City News.*

"A very businesslike and serviceable collection of essays and notes on this intricate question."—*Manchester Guardian.*

MANCHESTER UNIVERSITY PUBLICATIONS.
ECONOMIC SERIES.

(GARTSIDE REPORT, No. 4.)

No. VI. DYEING IN GERMANY AND AMERICA. By SYDNEY H. HIGGINS, M.Sc., Gartside Scholar. Demy 8vo, pp. xiii. 112. 1s. net. (Publication No. 24, 1907.)

"The book will . . . make a valuable addition to the technical literature of this country."—*Tribune.*

"The work is one which . . . . should receive the attention of those who desire a general view of the German and American dyeing industries."—*Textile Manufacturer.*

"A perusal of the work leads us to the conclusion that much useful work is being done by the Gartside scholars, which will give these young men an excellent insight into the working conditions of various industries."—*Textile Recorder.*

No. VII. THE HOUSING PROBLEM IN ENGLAND. By ERNEST RITSON DEWSNUP, M.A., Professor of Railway Economics in the University of Chicago. Demy 8vo, pp. vii. 327. 5s. net. (Publication No. 25, 1907.)

"Mr. Dewsnup's book is most valuable as it provides all essential information on the subject."—*Standard.*

"All those who are interested in this question, no matter what their economic predilections, may ponder with advantage Professor Dewsnup's pages."—*Newcastle Daily Chronicle.*

"The study brings together so weighty an array of facts and arguments that it cannot but prove instructive and suggestive to all classes of economists interested in its subject."—*Scotsman.*

"Professor Dewsnup's view of the whole problem was stated in 1903, in a form which won the Warburton Essay Prize at the Manchester University. Now revised and brought up to date, his valuable work has taken permanent form."—*Westminster Gazette.*

(GARTSIDE REPORT, No. 5.)

No. VIII. AMERICAN BUSINESS ENTERPRISE. By DOUGLAS KNOOP, M.A. Price 1s. 6d. net. (Publication No. 30, 1907.)

"The book is calculated to give a clear and accurate description, "essentially intended for the general reader," and the author has quite rightly eliminated everything of a technical character, giving his theme both the simplicity and the interest that are required. . . . The work might well have been doubled in length without any loss of interest. . . . Invaluable as a text-book."—*The Economic Journal.*

"Should on no account be missed, for it is a very good attempt at a survey of the enormous field of American business in the true and judicial spirit."—*Pall Mall Gazette.*

"Readable, informing, suggestive—full of interest for men engaged in almost every department of commercial life."—*Manchester City News.*

"A report of the general conditions of industrial work in the United States, together with a most instructive review of the education of the business man in their commercial universities."

—*Manchester Daily Dispatch.*

"The report is full of information, and is suggestive throughout."

—*Liverpool Post.*

"Concise, business-like and informative, it emphasises the difference between the economic positions of England and of America, and cannot but prove instructive to anyone interested in its subject."—*Scotsman.*

"From the point of view of an intelligent observer and collator, trained, alert, well-informed, bringing his mind to bear on the fundamental elements of commercial progress and success, it would be impossible to estimate it too highly."—*Belfast Northern Whig.*

No. IX. THE ARGENTINE AS A MARKET. By N. L. WATSON. Demy 8vo. 1s. net. (Publication No. 33, 1908.)

MANCHESTER UNIVERSITY PUBLICATIONS.

EDUCATIONAL SERIES.

No. I. CONTINUATION SCHOOLS IN ENGLAND & ELSEWHERE. Their place in the Educational System of an Industrial and Commercial State. By MICHAEL E. SADLER, M.A., LL.D., Professor of the History and Administration of Education. Demy 8vo, pp. xxvi. 779. 8s. 6d. net. (Publication No. 29, 1907.)

This work is largely based on an enquiry made by past and present Students of the Educational Department of the University of Manchester. Chapters on Continuation Schools in the German Empire, Switzerland, Denmark, and France, have been contributed by other writers.

". . . . gives a record of what the principal nations are doing in the prolongation of school work. It is invaluable as a *corpus* of material from which to estimate the present position of the world—so far as its analogies touch Britain—in 'further education,' as the phrase is."
—*The Outlook.*

"The most comprehensive book on continuation schools that has yet been issued in this country."—*Scottish Review.*

"Professor Sadler has produced an admirable survey of the past history and present condition of the problem of further education of the people . . . . but apart from his own contributions, the bulk of the work, and its most valuable portion, consists of material furnished by teachers and by organisers of schools in various parts of England and Scotland, by officials of the Board of Education and the Board of Trade, and by local education authorities."—*Manchester Guardian.*

"A perfect mine of facts and opinions. . . . is certain of a hearty welcome from all engaged in administering education."—*Glasgow Herald.*

"This is a book which counts. It is a worthy treatment of an all-important subject, and he who wishes his country well must pray that it may be read widely. . . . . I should be glad to think that I have said enough to send many readers post-haste to buy this invaluable treatise."
—L. J. Chiozza Money, M.P., in the *Daily News.*

"Professor Sadler's book is an admirable work on a subject which has not hitherto been dealt with in so masterly and complete a manner."
—*Manchester City News.*

"A volume which may mark a new epoch in educational thought and effort in England."—*The Tribune.*

"This book will for many years remain the standard authority upon its subject."—*The Guardian.*

"It is indeed a remarkable compilation, and we hope that its circulation and its usefulness may be commensurable with its conspicuous merits."—*The Schoolmaster.*

"The whole question is discussed with an elaboration, an insistence on detail, and a wisdom that mark this volume as the most important contribution to educational effort that has yet been made."
—*Contemporary Review.*

"This is a most valuable and opportune book, one to be commended to the careful attention of every serious student of the social problem."
—*The Churchman.*

"The book brims with interest to every man who recognizes the need of greater educational ideals in the masses."—*Co-operative News.*

"A work which we strongly recommend to all interested in the study of the social problem."—*The Record.*

MANCHESTER UNIVERSITY PUBLICATIONS.

EDUCATIONAL SERIES.

"The subject of the work is one that goes to the very heart of national education, and the treatise itself lays bare with a scientific but humane hand the evils that beset our educational system, the waste of life and national energy which that system has been unable in any sufficient degree to check."—*The Spectator.*

"It is a treasure of facts and judicious opinions in the domain of the history and administration of education."—*The Athenæum.*

"The volume represents an immense service to English education, and to the future welfare and efficiency of the nation."—*Educational Times.*

No. II. THE DEMONSTRATION SCHOOLS RECORD. No. I. Being Contributions to the Study of Education from the Department of Education in the University of Manchester. By Professor J. J. FINDLAY. 1s. 6d. net. (Publication No. 32, 1908.)

"This volume marks a new departure in English Educational literature . . . . Some very interesting work is being done and the most valuable part of the book is the account of the detailed methods which have been employed both in the regular teaching in the schools and in the efforts to foster the corporate interests of the children and their parents. These methods are often exceedingly suggestive, and may be studied with advantage by those who do not accept all the theories upon which they are based."—*School.*

"Professor Findlay and his skilled and experienced collaborators give an interesting account of the uses of the demonstration classes, the nature and scope of the work done in them, and the methods adopted (as well as the underlying principles) in some of the courses of instruction."—*The Athenæum.*

"The book gives an instructive account of the attempts made to correlate the subjects of school instruction, not only with each other, but also with the children's pursuits out of school hours. . . . The problem Professor Findlay has set himself to work out in the Demonstration School is, How far is it possible by working with the children through successive culture epochs of the human race to form within their minds not only a truer conception of human history, but also eventually a deeper comprehension of the underlying purpose and oneness of all human activities?"—*Morning Post.*

"Here the authors take us into their confidence; we are told what their view of a demonstration school is, what questions they hope to solve, and on what principles they think the answers should be sought. . . . . Those interested in educational progress will give the volume a cordial welcome."—*Nature.*

No. III. THE TEACHING OF HISTORY IN GIRLS' SCHOOLS IN NORTH AND CENTRAL GERMANY. A Report by EVA DODGE, M.A. (Publication No. 34, 1908.)

HISTORICAL SERIES.

No. I. MEDIÆVAL MANCHESTER AND THE BEGINNINGS OF LANCASHIRE. By JAMES TAIT, M.A., Professor of Ancient and Mediæval History. Demy 8vo, pp. x. 211. 7s. 6d. net. (Publication No. 3, 1904.)

"Patient and enlightened scholarship and a sense of style and proportion have enabled the writer to produce a work at once solid and readable."—*English Historical Review.*

"A welcome addition to the literature of English local history, not merely because it adds much to our knowledge of Manchester and Lancashire, but also because it displays a scientific method of treatment

MANCHESTER UNIVERSITY PUBLICATIONS.

HISTORICAL SERIES.

which is rare in this field of study in England."—Dr. Gross in *American Historical Review*.

"La collection ne pouvait débuter plus significativement et plus heureusement que par un ouvrage d'histoire du Moyen Age dû à M. Tait, car l'enseignement mediéviste est un de ceux qui font le plus d'honneur à la jeune Université de Manchester, et c'est à M. le Professeur Tait qu'il faut attribuer une bonne part de ce succès."—*Revue de Synthèse historique*.

"The two essays are models of their kind."—*Manchester Guardian*.

No. II. INITIA OPERUM LATINORUM QUAE SAECULIS XIII., XIV., XV. ATTRIBUUNTUR. By A. G. Little, M.A., Lecturer in Palæography. Demy 8vo, pp. xiii. 273 (interleaved). 15s. net. (Publication No. 5, 1904.)

"Whoever has attempted to ascertain the contents of a Mediæval miscellany in manuscript must often have been annoyed by the occurrence of a blank space where the title of the treatise ought to be. Mr. Little has therefore earned the gratitude of all such persons by making public a collection of some 6,000 incipits, which he arranged in the first instance for his private use, in compiling a catalogue of Franciscan MSS."
—*English Historical Review*.

No. III. THE OLD COLONIAL SYSTEM. By Gerald Berkeley Hertz, M.A., B.C.L., Lecturer in Constitutional Law. Demy 8vo, pp. xi. 232. 5s. net. (Publication No. 7, 1905.)

"Mr. Hertz gives us an elaborate historical study of the old colonial system, which disappeared with the American Revolution. . . . . He shows a remarkable knowledge of contemporary literature, and his book may claim to be a true history of popular opinion."—*Spectator*.

"Mr. Hertz's book is one which no student of imperial developments can neglect. It is lucid, fair, thorough, and convincing."
—*Glasgow Herald*.

"Mr. Hertz's 'Old Colonial System' is based on a careful study of contemporary documents, with the result that several points of no small importance are put in a new light . . . . it is careful, honest work . . . . The story which he tells has its lesson for us."—*The Times*.

"Both the ordinary reader and the academic mind will get benefit from this well-informed and well-written book."—*Scotsman*.

"Mr. Hertz has made excellent use of contemporary literature, and has given us a very valuable and thorough critique. The book is interesting and very well written."—*American Political Science Review*.

"An interesting, valuable, and very necessary exposition of the principles underlying the colonial policy of the eighteenth century."
—*Yorkshire Post*.

"A work embodying much work and research. . . . Three most impressive chapters should be read by everyone."—*Birmingham Post*.

"Very enlightening."—*American Historical Review*.

"Timely and useful."—*Athenæum*.

No. IV. STUDIES OF ROMAN IMPERIALISM. By W. T. Arnold, M.A. Edited by Edward Fiddes, M.A., Lecturer in Ancient History, with Memoir of the Author by Mrs. Humphry Ward and C. E. Montague. With a Photogravure of W. T. Arnold. Demy 8vo, 400 pp. 7s. 6d. net. (Publication No. 16, 1906.)

"Mrs. Humphry Ward has used all her delicate and subtle art to draw a picture of her beloved brother; and his friend Mr. Montague's account of his middle life is also remarkable for its literary excellence."
—*Athenæum*.

MANCHESTER UNIVERSITY PUBLICATIONS.

HISTORICAL SERIES.

"The memoir . . . . . tenderly and skilfully written by the 'sister and friend,' tells a story, which well deserved to be told, of a life rich in aspirations, interests, and friendships, and not without its measure of actual achievement."—*Tribune.*

"This geographical sense and his feeling for politics give colour to all he wrote."—*Times.*

"Anyone who desires a general account of the Empire under Augustus which is freshly and clearly written and based on wide reading will find it here."—*Manchester Guardian.*

"Nothing could be better than the sympathetic tribute which Mrs. Humphry Ward pays to her brother, or the analysis of his work and method by his colleague Mr. Montague. The two together have more stuff in them than many big books of recent biography."

—*Westminster Gazette.*

The Memoir may be had separately, price 2s. 6d. net.

No. V. CANON PIETRO CASOLA'S PILGRIMAGE TO JERUSALEM IN THE YEAR 1494. By M. M. Newett, B.A., formerly Jones Fellow. Demy 8vo, pp. 427. 7s. 6d. net.
(Publication No. 26, 1907.)

"Thoroughness is characteristic of introduction, the copious notes, appendix and index. . . . Miss Newett's translation is spirited and interesting. . . ."—*Manchester Courier.*

"Casola's narrative richly deserved the honours of print and translation. The book is a credit to its editor and to the historical school of Manchester University."—*Morning Leader.*

"His narrative is at once simple and dignified in style, convincing and interesting in its pictures of the conditions governing travel by sea and land four centuries ago."—*Daily Telegraph.*

"The book is like a gallery of mediæval paintings, full of movement and colouring, instinct with the vitality of the time."—*Birmingham Post.*

"Miss Newett's introduction is a contribution of considerable value to the history of European commerce."—*Spectator.*

"Forms a noteworthy addition to the number of books from which a knowledge can be gained of the itineraries of the pilgrims to Palestine."

—*Scotsman.*

"The whole volume is fascinating. It presents a lively picture of bygone times, abounds in curious facts and recalls quaint and pleasing ceremonies, and exhibits the ardent pilgrim of the past in his true light. Miss Newett is alike to be congratulated on her translation, her Introduction (which takes up a third of the volume), and her notes."

—*Manchester City News.*

"The work which Miss Margaret Newett has probably saved from oblivion is as intrinsically interesting as it should prove instructive to the student of history."—*Daily News.*

"One of the most delightful narratives that record the impressions of a pious pilgrim."—*Westminster Gazette.*

"One of the most comprehensive of the itineraries is that now translated, an important feature of it being its full description of the city of Venice."—*The Times.*

No. VI. HISTORICAL ESSAYS. Edited by T. F. Tout, M.A., Professor of Mediæval and Modern History, and James Tait, M.A., Professor of Ancient and Mediæval History. Demy 8vo, pp. xv. 557. 6s. net. Reissue of the Edition of 1902 with Index and New Preface.
(Publication No. 27, 1907.)

"Diese zwanzig chronologisch geordneten Aufsätze heissen in der Vorrede der Herausgeber *Festschrift*, behandeln zur Hälfte ausser-englische

MANCHESTER UNIVERSITY PUBLICATIONS.

HISTORICAL SERIES.

Themata, benutzen reichlich festländische Literatur und verraten überall neben weiten Ausblicken eine methodische Schulung die der dortigen Facultät hohe Ehre macht."—Professor Liebermann in *Deutsche Literaturzeitung*.

"Imperial history, local history, ecclesiastical history, economic history and the methods of historical teaching—all these are in one way or another touched upon by scholars who have collaborated in this volume. Men and women alike have devoted their time and pains to working out problems of importance and often of no slight difficulty. The result is one of which the university and city may be justly proud."—The late Professor York Powell in the *Manchester Guardian*.

"Esso contiene venti lavori storici dettati, quattro da professori e sedici da licenziati del Collegio, e sono tutto scritti appositamente e condotti secondo le più rigorose norme della critica e su documenti."—R. Predelli in *Nuovo Archivio Veneto*.

"La variété des sujets et l'érudition avec laquelle ils sont traités font grand honneur à la manière dont l'histoire est enseigné à Owens College."
—*Revue Historique*.

"No one who reads these essays will do so without acknowledging their ability, both in originality and research. They deal with historic subjects from the beginnings of Cæsar-worship to the detention of Napoleon at St. Helena, and they deal with them in a thoroughgoing fashion."—*Guardian*.

"Par nature, c'est un recueil savant, qui témoigne du respect et de l'émulation que sait exercer pour les études historiques la jeune et déjà célèbre université."—*Revue d'histoire ecclésiastique* (Louvain).

"All these essays reach a high level; they avoid the besetting sin of most of our present historical writing, which consists of serving up a hash of what other historians have written flavoured with an original spice of error. . . . . They are all based on original research and written by specialists."—Professor A. F. Pollard in the *English Historical Review*.

"Sie bilden einen schönen Beweis fur die rationelle Art, mit der dort dieses Studium betrieben wird."—Professor O. Weber in *Historische Zeitschrift*.

The Index can be purchased separately, price 6d.

MEDICAL SERIES.

No. I. SKETCHES OF THE LIVES AND WORK OF THE HONORARY MEDICAL STAFF OF THE ROYAL INFIRMARY. From its foundation in 1752 to 1830, when it became the Royal Infirmary. By Edward Mansfield Brockbank, M.D., M.R.C.P. Crown 4to. (illustrated). Pp. vii. 311. 15s. net.
(Publication No. 1, 1904.)

"Dr. Brockbank's is a book of varied interest. It also deserves a welcome as one of the earliest of the 'Publications of the University of Manchester.'"—*Manchester Guardian*.

"We have a valuable contribution to local Medical Liteurature."
—*Daily Dispatch*.

No. II. PRACTICAL PRESCRIBING AND DISPENSING. For Medical Students. By William Kirkby, sometime Lecturer in Pharmacognosy in the Owens College, Manchester. Crown 8vo, 220 pp. 5s. net.
(Publication No. 2, 1904, Second edition, 1906.)

"The whole of the matter bears the impress of that technical skill and thoroughness with which Mr. Kirkby's name must invariably be

MANCHESTER UNIVERSITY PUBLICATIONS.

MEDICAL SERIES.

associated, and the book must be welcomed as one of the most useful recent additions to the working library of prescribers and dispensers."
—*Pharmaceutical Journal.*

"Thoroughly practical text-books on the subject are so rare, that we welcome with pleasure Mr. William Kirkby's 'Practical Prescribing and Dispensing.' The book is written by a pharmacist expressly for medical students, and the author has been most happy in conceiving its scope and arrangement."—*British Medical Journal.*

"The work appears to be peculiarly free from blemishes and particularly full in practical detail. It is manifestly the work of one who is a skilled chemist, and an expert pharmacist, and who knows not only the requirements of the modern student but the best way in which his needs may be met."—*Medical Press.*

"This is a very sensible and useful manual."—*The Hospital.*

"The book will be found very useful to any students during a course of practical dispensing."—*St Bartholomew's Hospital Journal.*

"The book is a model, being tutorial from beginning to end."
—*The Chemist and Druggist.*

No. III. HANDBOOK OF SURGICAL ANATOMY. By G. A. Wright, B.A., M.B. (Oxon.), F.R.C.S., Professor of Systematic Surgery, and C. H. Preston, M.D., F.R.C.S., L.D.S., Lecturer on Dental Anatomy; Assistant Dental Surgeon to the Victoria Dental Hospital of Manchester. Crown 8vo, pp. ix. 205. Second edition. 5s. net. (Publication No. 6, 1905.)

"We can heartily recommend the volume to students, and especially to those preparing for a final examination in surgery."—*Hospital.*

"Dr. Wright and Dr. Preston have produced a concise and very readable little handbook of surgical applied anatomy. . . . The subject matter of the book is well arranged and the marginal notes in bold type facilitate reference to any desired point."—*Lancet.*

No. IV. A COURSE OF INSTRUCTION IN OPERATIVE SURGERY in the University of Manchester. By William Thorburn, M.D., B.S. (Lond.), F.R.C.S., Lecturer in Operative Surgery. Crown 8vo, pp. 75. 2s. 6d. net.
(Publication No. 11, 1906.)

"This little book gives the junior student all that he wants, and nothing that he does not want. Its size is handy, and altogether for its its purpose it is excellent."—*University Review.*

"As a working guide it is excellent."—*Edinburgh Medical Journal.*

No. V. A HANDBOOK OF LEGAL MEDICINE. By W. Sellers, M.D. (London), of the Middle Temple, and Northern Circuit, Barrister-at-law. With Illustrations. Crown 8vo, pp. vii. 233. 7s. 6d. net. (Publication No. 14, 1906.)

"This is quite one of the best books of the kind we have come across."—*Law Times.*

No. VI. A CATALOGUE OF THE PATHOLOGICAL MUSEUM OF THE UNIVERSITY OF MANCHESTER. Edited by J. Lorrain Smith, M.A., M.D. (Edin.), Professor of Pathology. Crown 4to, 1260 pp. 7s. 6d. net. (Publication No. 15, 1906.)

"The catalogue compares very favourably with others of a similar character, and, apart from its value for teaching purposes in an important medical school such as that of the University of Manchester, it is capable of being of great assistance to others as a work of reference."

MANCHESTER UNIVERSITY PUBLICATIONS.

MEDICAL SERIES.

—*Edinburgh Medical Journal.*

"In conclusion we need only say that Professor Lorrain Smith has erformed the most essential part of his task—the description of the ecimens—excellently, and an honourable mention must be made of e book as a publication"—*British Medical Journal.*

o. VII. HANDBOOK OF DISEASES OF THE HEART. By Graham Steell, M.D., F.R.C.P., Professor of Medicine, and Physician to the Manchester Royal Infirmary. Crown 8vo, pp. xii. 389, 11 plates (5 in colours), and 100 illustrations in the text. 7s. 6d. net. (Publication No. 20, 1906.)

"It more truly reflects modern ideas of heart disease than any book e are acquainted with, and therefore may be heartily recommended to ur readers."—*Treatment.*

"We regard this volume as an extremely useful guide to the study of iseases of the heart, and consider that no better introduction to the ubject could possibly have been written."

—*Medical Times and Hospital Gazette.*

"We can cordially recommend Dr. Steell's book as giving an excellent nd thoroughly practical account of the subject of which it treats."—*Edinburgh Medical Review.*

o. VIII. JULIUS DRESCHFELD. IN MEMORIAM. Medical Studies by his colleagues and pupils at the Manchester University and the Royal Infirmary. (Publication No. 35, 1908.)

PHYSICAL SERIES.

o. I. THE PHYSICAL LABORATORIES OF THE UNIVERSITY OF MANCHESTER. A record of 25 years' work. Demy 8vo, pp. 142, 10 Plates, 4 Plans. 5s. net. (Publication No. 13, 1906.)

This volume contains an illustrated description of the Physical, Electrical Engineering, and Electro-Chemistry Laboratories of the Manchester University, also a complete Biographical and Bibliographical Record of those who have worked in the Physics Department of the University during the past 25 years.

"The book is excellently got up, and contains a description of the epartment of physics and its equipment, a short biographical sketch of he Professor with a list of his scientific writings and a well-executed ortrait and a record of the career of students and others who have passed hrough Dr. Schuster's hands. Alumni of Owens will welcome the olume as an interesting link with their alma mater."—*Glasgow Herald.*

"This interesting and valuable contribution to the history of the Manchester University also contains several illustrations, and forms the irst of the 'physical series' of the publications of the University of Manchester."—*The Times.*

"A record of achievement of which no man need be ashamed."

—*Westminster Gazette.*

"It is a memorial of which any man would be justly proud, and the University of which he is both an alumnus and a professor may well share that pride."—*Manchester Guardian.*

PUBLIC HEALTH SERIES.

No. I. ARCHIVES OF THE PUBLIC HEALTH LABORYTORY OF THE UNIVERSITY OF MANCHESTER. Edited by A. Sheridan Delépine, M.Sc., M.B., Ch. M., Director of the Laboratory and Proctor Professor of Comparative Pathology and Bacteriology. Crown 4to. pp. iv. 451. £1. 1s. net. (Publication No. 12, 1906.)

MANCHESTER UNIVERSITY PUBLICATIONS.

PUBLIC HEALTH SERIES.

"The University of Manchester has taken the important and highly commendable step of commencing the publication of the archives of its Public Health Laboratory, and has issued, under the able and judicious editorship of Professor Sheridan Delépine, the first volume of a series that promises to be of no small interest and value alike to members of the medical profession and to those of the laity. . . . Original communications bearing upon diseases which are prevalent in the districts surrounding Manchester, or dealing with food- and water-supplies, air, disposal of refuse, sterilisation and disinfection and kindred subjects, will be published in future volumes; and it is manifest that these, as they successively appear, will form a constantly increasing body of trustworthy information upon subjects which are not only of the highest interest to the profession but of supreme importance to the public."
—*The Lancet.*

"It is safe to say that as these volumes accumulate they will form one of the most important works of reference on questions of public health, and ought, at all events, to be in the library of every public authority."—*Manchester Guardian.*

"The volume . . . . speaks well for the activity of investigation in Manchester."—*Lancet.*

THEOLOGICAL SERIES.

No. I. INAUGURAL LECTURES delivered during the Session 1904–5, by the Professors and Lecturers of the Faculty of Theology, viz. :—

Prof. T. F. Tout, M.A.; Prof. A. S. Peake, B.D.; Prof. H. W. Hogg, M.A.; Prof. T. W. Rhys Davids, LL.D.; Rev. W. F. Adeney, D.D.; Rev. A. Gordon, M.A.; Rev. L. Hassé, B.D.; Rev. Canon E. L. Hicks, M.A.; Rev. H. D. Lockett, M.A.; Rev. R. Mackintosh, D.D.; Rev. J. T. Marshall, D.D.; Rev. J. H. Moulton, D.Litt.

Edited by A S. Peake, B.D., Dean of the Faculty.
Demy 8vo, pp. xi. 296. 7s. 6d. net.
(Publication No. 9, 1905.)

"The lectures, while scholarly, are at the same time popular, and will be found interesting and instructive by those who are not theologians. . . . . The entire series is excellent, and the volume deserves a wide circulation."—*Scotsman.*

"This is a very welcome volume. . . . All these lectures were delivered to popular audiences, yet they are far from superficial, and will be found of great value to busy pastors and teachers."—*Christian World.*

"We welcome the volume as a most auspicious sign of the times."
—*Spectator.*

"The lectures themselves give a valuable conspectus of the present position of Theological research. . . . They are, of course, not addressed to experts, but they are exceedingly valuable, even when allowance is made for their more or less popular form."—*Examiner.*

"The whole volume forms a very important and valuable contribution to the cause of Theological learning."—*Record.*

"This is a most interesting and valuable book, the appearance of which at the present moment is singularly significant. . . . But it is impossible in a brief review to indicate all the treasures of this rich volume, to read which carefully is to be introduced to the varied wealth of modern Biblical scholarship."—*Baptist.*

"This volume is of the most exceptional value and interest."
—*Expository Times.*

MANCHESTER UNIVERSITY PUBLICATIONS.

THEOLOGICAL SERIES.

"This is a book of more than common interest."
—*Review of Theology and Philosophy.*

"The writers of these lectures do not attempt to offer more than mples of their wares: but what is given is good, and it may be seen at theology without tests is destitute neither of scientific value nor of ıman interests."—*Athenæum.*

LECTURES.

o. I. GARDEN CITIES (Warburton Lecture). By RALPH NEVILLE, K.C. 6d. net. (Lecture No. 1, 1905.)

o. II. THE BANK OF ENGLAND AND THE STATE (A Lecture). By Sir FELIX SCHUSTER. 6d. net. (Lecture No. 2, 1905.)

o. III. BEARING AND IMPORTANCE OF COMMERCIAL TREATIES IN THE TWENTIETH CENTURY. By Sir THOMAS BARCLAY. 6d. net. (Lecture No. 3, 1906.)

o. IV. THE SCIENCE OF LANGUAGE AND THE STUDY OF THE GREEK TESTAMENT (A Lecture). By JAMES HOPE MOULTON, M.A., Litt.D. 6d. net. (Lecture No. 4, 1906.)

o. V. THE GENERAL MEDICAL COUNCIL: ITS POWERS AND ITS WORK (A Lecture). By DONALD MACALISTER, M.A., M.D., B.Sc., D.C.L., LL.D. 6d. net. (Lecture No. 5, 1906.)

No. VI. THE CONTRASTS IN DANTE (A Lecture). By the Hon. WILLIAM WARREN VERNON, M.A. 6d. net. (Lecture No. 6, 1906.)

No. VII. THE PRESERVATION OF PLACES OF INTEREST OR BEAUTY (A Lecture). By SIR ROBERT HUNTER. 6d. net. (Lecture No. 7, 1907.)

CALENDARS

CALENDAR OF THE VICTORIA UNIVERSITY OF MANCHESTER. Session 1904-5. Demy 8vo, 1100 pp. 3s. net. (Publication No. 17.)

CALENDAR OF THE VICTORIA UNIVERSITY OF MANCHESTER. Session 1905-6. Demy 8vo, 1200 pp. 3s. net. (Publication No. 18.)

CALENDAR OF THE VICTORIA UNIVERSITY OF MANCHESTER. Session 1906-7. Demy 8vo, 1300 pp. 3s. net. (Publication No. 19.)

CALENDAR OF THE VICTORIA UNIVERSITY OF MANCHESTER. Session 1907-8. Demy 8vo, 1400 pp. 3s. net. (Publication No. 28.)

CALENDAR OF THE VICTORIA UNIVERSITY OF MANCHESTER. Session 1908-9. Demy 8vo, 1460 pp. 3s. net. (Publication No. 37.)

THE REGISTER OF GRADUATES OF THE UNIVERSITY OF MANCHESTER UP TO JULY 1908. 2s. 6d. net, cloth 3s. 6d. net.

MANCHESTER UNIVERSITY PUBLICATIONS.

Celtic Series. No. I.

The following are in preparation and will be issued shortly :—

AN INTRODUCTION TO EARLY WELSH. By the late Prof. J. Strachan, M.A., LL.D. Edited and completed by Prof. Kuno Meyer, Ph.D. Demy 8vo.

This work will comprise a Grammar of Early Welsh with special reference to Middle-Welsh prose. To the grammar will be added selected passages from Early Welsh texts in prose and verse, together with notes and a glossary compiled by Timothy Lewis, B.A. *[In October.*

A GLOSSARY TO THE BLACK BOOK OF CHIRK MANUSCRIPT OF THE WELSH LAWS. By Timothy Lewis, B.A. Demy 8vo.

This will include a complete glossary to the oldest copy of the "Laws of Howel Dda," contained in the "Black Book of Chirk," and will be based on the photographic facsimile of that manuscript which is about to be published by Dr. J. Gwenogvryn Evans in his collection of Welsh texts. *[In Preparation.*

THE LANGUAGE OF THE ANNALS OF ULSTER. By Tomás O'Máille, M.A. Demy 8vo.

The objects of this dissertation are firstly to investigate the date at which certain old-Irish phonological developments took place, and secondly to give an account of old-Irish declension as evidenced by the language of the Annals of Ulster. An Appendix on the analysis of Irish personal names is appended. *[In Preparation.*

Economic Series.

SOME ELECTRO-CHEMICAL CENTRES. Gartside Report. By J. N. Pring, M.Sc. *[In the Press.*

Historical Series.

STUDIES SUPPLEMENTARY TO STUBBS' CONSTITUTIONAL HISTORY. Vol. i. By Ch. Petit-Dutaillis, Litt.D., rector of the University of Grenoble. Translated from the French by W. E. Rhodes, M.A., and edited by Prof. James Tait, M.A.

This work will consist of the translation of the studies and notes appended by Prof. Petit-Dutaillis to his translation into French of the first volume of Stubbs' *Constitutional History of England.* It is believed that they will present to English students and teachers a summary of the results of recent historical research so far as they throw light upon or modify the conclusions expressed thirty years ago by the late Bishop Stubbs. *[In October.*

HANES GRUFFYDD AP CYNAN. The Welsh text with translation, introduction, and notes by Arthur Jones, M.A., Jones Fellow in History. Demy 8vo. *[In Preparation.*

SHERRATT & HUGHES

MANCHESTER UNIVERSITY PUBLICATIONS.

HE CROMWELLIAN CONQUEST AND SETTLEMENT OF IRELAND. By Robert Dunlop, M.A., formerly Berkeley Fellow. Demy 8vo.

This work will consist of a series of unpublished documents relating the History of Ireland from 1651 to 1659, arranged, modernized, and lited, with introduction, notes, etc., by Mr. Dunlop.
[*In Preparation.*

Medical Series.

ANDBOOK OF INFECTIOUS DISEASES. By R. W. Marsden, M.D. [*Immediately.*

ODERN PROBLEMS IN PSYCHIATRY. By E. Lugaro, Professor of Nervous and Mental Diseases in the University of Modena. Translated from the Italian by David Orr, M.D., Assistant Medical Officer and Pathologist to the County Asylum, Prestwich; and R. G. Rows, M.D., Assistant Medical Officer and Pathologist to the County Asylum, Lancaster. With an introduction by T. S. Clouston, M.D., Physician Superintendent, Royal Asylum, Morningside, and Lecturer on Mental Diseases in Edinburgh University.

Deals with the problems met with in studying the causation of innity. These problems are discussed under the headings of psychogical, anatomical, pathogenetic, etiological, nosological, social and actical. There are 13 illustrations in the anatomical section.
[*In Preparation.*

ISEASES OF THE EAR. By W. Milligan, M.D., Lecturer on Diseases of the Ear and Nasal Surgeon to the Manchester Royal Infirmary. [*In Preparation.*

ISEASES OF THE EYE. By C. E. Glascott, M.D., Lecturer on Ophthalmology, and A. Hill Griffith, M.D., Ophthalmic Surgeon to the Manchester Royal Infirmary. *In Preparation.*

Zoological Series.

TRUCTURE, DEVELOPMENT AND BIONOMICS OF THE HOUSE FLY. By C. Gordon Hewitt, M.Sc. [*In Preparation.*

The following works, though not technically Publicaions of the University of Manchester, are also issued from he University Press :—

IELANDRA CASTLE, being the Report of the Manchester and District Branch of the Classical Association for 1905. Edited by R. S. Conway, Litt.D. Introduction by Rev. E. L. Hicks, M.A. Demy 8vo. Illustrated. 5s. net.

RANSACTIONS OF THE INTERNATIONAL UNION FOR CO-OPERATION IN SOLAR RESEARCH (Vol. i., First and Second Conferences). Demy 8vo, 260 pp. and plate. 7s. 6d. net.

HE BOOK OF RUTH (Unpointed Text). 6d. net.

HE BOOK OF AMOS. (Unpointed Text.) 6d. net.

60, Chandos Street, London, W.C.

SHERRATT & HUGHES

MANCHESTER UNIVERSITY PUBLICATIONS.

SCENES FROM THE RUDENS OF PLAUTUS, with a Translation into English Verse. Edited by R. S. Conway, Litt.D., Professor of Latin in the University. 6d. net.

THE MOSTELLARIA OF PLAUTUS. Acting edition with a translation into English Verse. Edited by G. Norwood, M.A. 1s. net.

THE TEACHING OF HISTORY AND OTHER PAPERS. By H. L. Withers. Edited by J. H. Fowler. Crown 8vo, 270 pp. 4s. 6d. net.

"An interesting memorial of a teacher who was a real enthusiast for education."—*The Times.*

"We can cordially commend this little book to the somewhat limited but slowly widening circle who are likely to be interested in educational principles and organization."—*The Guardian.*

A TARDINESS IN NATURE AND OTHER PAPERS. By Mary Christie. Edited, with Introductory Note and Memoir, by Maud Withers. Crown 8vo, 331 pp. 3s. net.

"The essays upon Thackeray, George Eliot, and R. L. Stevenson in this volume could scarcely be bettered."—*The Guardian.*

"The life-story of a quite remarkable woman—of a woman who used her gifts always to the furthering of all that is sweetest and noblest in life."—*Tribune.*

MUSICAL CRITICISMS. By Arthur Johnstone. With a Memoir of the Author by Henry Reece and Oliver Elton. Crown 8vo, 225 pp. 5s. net.

"Without the smallest affectation or laboured attempts at smartness, Mr. Johnstone contrived always to throw fresh light on the matter in hand, and at the same time to present his opinions in a form which could be understood and enjoyed by the non-musical reader."

—*Westminster Gazette.*

"Everyone who welcomes guidance as to what is best in music, everyone who watches with some degree of fascination the power of analysis, everyone who reads with a sense of satisfaction English, as it may be written by a master of the craft, should read this book."

—*The Musical World.*

MANCHESTER BOYS. By C. E. B. Russell. With an Introduction by E. T. Campagnac. Crown 8vo. 2s. 6d. net.

"Mr. Charles E. B. Russell has written a most interesting and thought-compelling book on a subject of almost vital importance."

—*Yorkshire Post.*

"Altogether it is an inspiring book."

—*Liverpool Daily Post and Mercury.*